

UDC 575.21: 631.523:
Original scientific paper

COMPONENTS OF PHENOTYPIC VARIABILITY FOR HEAD DIAMETER IN SUNFLOWER (*HELIANTHUS ANNUUS* L.)

Nada HLADNI, Dragan ŠKORIĆ, and Marija KRALJEVIĆ-BALALIĆ¹

Institute of Field and Vegetable Crops, 21000 Novi Sad

¹ Faculty of Agriculture, 21000 Novi Sad, Serbia and Montenegro

Hladni N., D. Škorić, and M. Kraljević-Balalić (2003):
Components of phenotypic variability for head diameter in sunflower
(*Helianthus annuus* L.). – Genetika, Vol. 35, No. 2, 67-75.

Sunflower is the main crop species for the production of edible oil in many countries of the world, including ours. Plant height and head size, form and position on the stem as well as the number of leaves, their size, duration and distribution on the plant all play an important role in defining optimal plant architecture in sunflower hybrids (ŠKORIĆ, 1975, 1989, 2002). In order to monitor the mode of inheritance and gene effects for head diameter in the F₁ and F₂ generations, half diallel crosses were made in six genetically divergent sunflower inbreds. The mean values of the trait in question differed significantly. The mode of inheritance of head diameter in the F₁ generation was superdominance in all but four crosses, which had dominance instead. In the F₂ generation, on the other hand, the prevailing mode was dominance, while superdominance was recorded in four cases. The dominant component accounted for the bulk of genetic variance, and the mode of inheritance of head diameter taking into account both the F₁ and F₂ generations was superdominance. These findings may prove valuable for developing high-yielding sunflower genotypes.

Key words: sunflower, inbred lines, mode of inheritance, gene effect, simple correlation coefficient

Corresponding author: Nada Hladni, Scientific Institute of Field and Vegetable Crops,
M.Gorkog 30, 21000 Novi Sad, Serbia and Montenegro

INTRODUCTION

The sunflower is the main oil crop in many countries including Serbia and Montenegro. This is why emphasis in breeding for yield is placed on high seed production per plant. This is achieved by developing hybrids that combine high genetic yield potential, desirable plant architecture, tolerance to diseases and adaptability to specific agroecological conditions.

ALBA *et al.* (1979), GIRARAJ *et al.* (1979) and MARINKOVIĆ (1987) reported significant direct effects of head diameter on seed yield. Indirect effects of head diameter on seed yield were reported by AHMAD *et al.* (1991).

Numerous researchers (ŠKORIĆ, 1975; IVANOV, 1980; MARINKOVIĆ, 1992; SUZER and ATAKISS, 1993; PETAKOV, 1994; PUNIA and GILL, 1994) have found significant positive correlations between morphological characteristics such as head diameter and seed yield.

By changing the hybrid model it is possible to increase the competitiveness of sunflower plants. Such plants will be able to produce more than 1,500 seeds per head when grown in a dense stand, i.e., they will be able to achieve high and stable seed yields and oil contents. A yield component that directly changes hybrid model is head size which is expressed as head diameter in cm. Head size should be intermediate, with a diameter of 20-25 cm, thin and with firm epidermis underside (ŠKORIĆ, 1980). Increase in head size above the optimum value results in reduced kernel yield (g/head), increased husk percentage, increased number of empty seeds and reduced oil content in seed (ŠKORIĆ, 1989).

In addition to genetic factors, head size is also affected by environmental factors (soil moisture and fertility, number of plants per unit area) and the length of growing season (MARINKOVIĆ *et al.*, 2003). FICK (1978) concluded that the contribution of genetic factors in the inheritance of head diameter is lower than for most of the other agronomic character.

Significant effect of plant population on head diameter ($r=-0.89^{**}$) was reported by MICHELE *et al.* (1996). Increased row distance tends to increase head diameter (ESENDAL and KANDEMIR, 1996).

Investigations on the inheritance of head diameter conducted so far have provided different results. The results varied in dependence of the material used for crossing. KOVAČIK and ŠKALOUD (1971), VOLF and DUMANČEVA (1973), SEETY *et al.* (1977), MARINKOVIĆ (1984), ŠEĆEROV-FIŠER (1992), KUMAR (1998) and JOKSIMOVIĆ *et al.* (2000) found that the non-additive genetic variance prevailed in the inheritance of head diameter. Conversely, PUTT (1965), KOVAČIK and ŠKALOUD (1972), RAO i SING (1977), SINDAGI *et al.* (1979), DUA *et al.* (1985), MARINKOVIĆ and ŠKORIĆ (1990) and EL-HITY (1992) reported that the additive component of genetic variance prevailed over the non-additive.

To efficiently use the available breeding material, it is necessary to have knowledge of its genetic makeup and of the mode of inheritance of the characteristic bred for (KRALJEVIĆ-BALALIĆ, 1986).

The objectives of this investigation were to assess the mode of inheritance of head diameter in the F₁ and F₂ generations of sunflower and to evaluate gene

effects and the relationship between head diameter and seed yield per sunflower plant

MATERIAL AND METHOD

To assess the mode of inheritance and gene effects for sunflower head diameter, six genetically divergent inbred lines (OCMS₁, NS-204B, NS-22B, NS-BD, NS-NDF, NS-K), developed at Institute of Field and Vegetable Crops in Novi Sad, were diallelly crossed, without reciprocals, to produce the F₁ and F₂ generations. The experiment was then established in the random blocks system in three replicates in the location of Rimski Šančevi. The experiment included the six inbred lines, 15 F₁ and 15 F₂ hybrids. Samples for analyses were taken from inner rows, excluding border rows. The samples of the parent lines and the F₁ generation consisted of five plants per replicate, of the F₂ generation of 30 plants per replicate. Measurements of the analyzed characteristic were done in the field, at maturity stage.

Mean values and correlation coefficient r , as an indicator of mutual interdependence between the two variables, were calculated according to HADŽIVUKOVIĆ (1991).

To assess the mode of inheritance for head diameter, we tested the significance of the mean values of the F₁ and F₂ generations in relation to the parents' mean (BOROJEVIĆ, 1965).

To assess gene effects for the studied characters, the combining ability of the diallel crosses according to GRIFFING (1956), method 2, model I were analyzed.

Genetic variances were assessed by the method of MATHER and JINKS (1982).

RESULTS AND DISCUSSION

The inbred lines and their F₁ and F₂ hybrids differed significantly in the mean values for head diameter and yield per plant (Table 1). The smallest diameter was found in line NS-NDF (16.3 cm), the largest in line OCMS₁ (21.2 cm). The lowest mean values in the F₁ and F₂ generations were found in combination NS-22BxNS-NDF (19.4 cm and 18.2 cm, respectively), the highest in combination OCMS₁xNS-BD (24.1 cm and 23.8 cm, respectively), Table 1.

In the F₁ generation, four combinations inherited head diameter dominantly from the parent with the larger mean value, while positive heterosis occurred in 11 combinations (Table 1).

In the F₂ generation, dominance of the parent with the larger mean value was registered in 11 combinations, and positive heterosis was exhibited in the following combinations: OCMS₁xNS-204B, OCMS₁xNS-BD, NS-22BxNS-K and NS-NDFxNS-K (Table 1). Dominance and superdominance was stated in the inheritance of head diameter (MOROZOV, 1947; SCHUSTER, 1964; KOVAČIK i ŠKALOUĐ, 1971; STOJANOVA *et al.*, 1971; MARINKOVIĆ, 1984).

Table 1. Mean values for head diameter (cm) and seed yield per plant

Parents and hybrids		Head diameter	Seed yield per plant
		cm	g
P ₁	OCMS ₁	21.2	43.3
F ₁	OCMS ₁ xNS204B	24.1 ^h	88.6
F ₂		23.3 ^h	81.6
F ₁	OCMS ₁ xNS22B	22.9 ^h	91.0
F ₂		21.9 ^{dt}	53.1
F ₁	OCMS ₁ xNSBD	24.1 ^h	100.7
F ₂		23.8 ^h	71.9
F ₁	OCMS ₁ xNSNDF	22.3 ^h	78.3
F ₂		21.5 ^{dt}	63.8
F ₁	OCMS ₁ xNSK	22.2 ^h	105.2
F ₂		21.5 ^{dt}	76.2
P ₂	NS204B	19.5	44.36
F ₁	NS204BxNS22B	21.9 ^h	102.5
F ₂		20.4 ^{dt}	51.4
F ₁	NS204BxNSBD	22.3 ^h	110.1
F ₂		20.8 ^{dt}	48.4
F ₁	NS204BxNSNDF	20.8 ^{dt}	81.3
F ₂		20.0 ^{dt}	48.4
F ₁	NS204BxNSK	21.2 ^h	109.7
F ₂		20.7 ^{dt}	65.2
P ₃	NS22B	18.2	47.2
F ₁	NS22BxNSBD	21.1 ^{dt}	90.2
F ₂		20.5 ^{dt}	56.8
F ₁	NS22BxNSNDF	19.4 ^{dt}	95.1
F ₂		18.2 ^{dt}	41.4
F ₁	NS22BxNSK	21.1 ^h	110.8
F ₂		20.0 ^h	70.0
P ₄	NSBD	20.7	43.6
F ₁	NSBDxNSNDF	22.3 ^h	92.2
F ₂		21.4 ^{dt}	56.8
F ₁	NSBDxNSK	22.5 ^h	128.9
F ₂		21.7 ^{dt}	76.3
P ₅	NSNDF	16.3	24.8
F ₁	NSNDFxNSK	20.8 ^h	115.4
F ₂		20.2 ^h	69.0
P ₆	NSK	17.4	29.7

The analysis of the components of genetic variance showed that the dominant component (H_1 and H_2) was larger than the additive one (D) for head diameter. It means that the dominant component makes the main portions of genetic variance in the F_1 and F_2 generations (Table 4). These results are in agreement with those of KOVAČIK and ŠKALOUD (1971), VOLF and DUMANČEVA (1973), SEETY *et al.* (1977), MARINKOVIĆ (1984) ŠEĆEROV-FIŠER (1992), KUMAR *et al.* (1998) and JOKSIMOVIĆ *et al.* (2000), but they are contrary to the results of PUTT (1965), KOVAČIK and ŠKALOUD (1972), RAO and SING (1977), SINDAGI *et al.* (1979), DUA *et al.* (1985), MARINKOVIĆ and ŠKORIĆ (1990) and EL HITY

(1992) who maintained that the additive component was more important than the dominant one in the inheritance of head diameter in sunflower.

As the F value was positive, it may be concluded that dominant genes prevail over recessive ones in the expression of this characteristic in the studied inbred lines. This was fully confirmed by the calculated gene frequencies. The frequency of dominant (u) genes was higher than the frequency of recessive (v) genes. Dominant (u) and recessive (v) genes were not evenly distributed among the parents as indicated by the ratio ($H_2/4H_1$) which was different from 0.25 (Table 2).

The calculated value of $(H_1/D)^{1/2}$, which shows the average degree of dominance, was larger than one. This was an indication that superdominance was exhibited in the inheritance of head diameter when all combinations in the F_1 and F_2 generations are taken into account (Table 2).

The ratio of the total numbers of dominant versus recessive genes for all parents showed that dominant genes prevailed since the values of Kd/Kr were larger than one in both generations (Table 2).

Table 2. Variance components for head diameter per sunflower plant

Component	Head diameter	
	Value	
	F_1	F_2
D	3.61	3.62
H_1	7.61	5.01
H_2	7.43	4.64
F	0.24	0.20
E	0.03	0.02
$H_2/4H_1$	0.24	0.23
u	0.58	0.65
v	0.42	0.35
$(H_1/D)^{1/2}$	1.45	1.18
Kd/Kr	1.05	1.05

The interdependence between head diameter and seed yield per plant was found to be positive and highly significant (0.601**), tab.3. That is in agreement with the results of (ŠKORIĆ, 1975; IVANOV, 1980; MARINKOVIĆ, 1992; SUZER and ATAKISS, 1993; PETAKOV, 1994; PUNIA and GILL, 1994).

Table 3. Correlation coefficient for head diameter and seed yield per plant

Characteristic	Head diameter	Seed yield per plant
Coefficient r		0.621**

** Significant at LSD 1%

This investigation may be useful in sunflower breeding for high genetic yield potential because it was found that the parents included into the diallel

possessed a large number of dominant genes for head diameter while highly significant interdependence was found between this parameter and seed yield per plant.

CONCLUSION

A diallel cross excluding reciprocals has been made in order to assess the mode of inheritance and gene effects for head diameter in the F_1 and F_2 generations of sunflower.

The mean values of the tested lines and hybrids differed significantly for the characteristics studied.

Head diameter was inherited dominant and superdominant in F_1 and F_2 .

The analysis of components of variance showed that dominant genes prevailed over recessive ones in the expression of head diameter, i.e., the major part of genetic variance was due to the dominant gene action.

The existence of superdominance in the inheritance of head diameter was confirmed by the average degree of dominance which was larger than unity.

This investigation may be important for further work because sunflower breeding for yield components based on heterosis should make use of crosses which exhibit non-additive gene effect (dominance and epistasis).

Received November 4th, 2003

Accepted December 29th, 2003

REFERENCES

- AHMAD A., M.A. RANA, and S.U.H. SIDIAQUI (1991): Sunflower seed yield as influenced by some agronomic and seed characters. *Euphytica*, 56, 137-142.
- ALBA E., A. BENVENUTI, R. TUBEROSA, and G.O. VANOZZI (1979): A path coefficient analysis of some yield components in sunflower. *Helia*, 2, 25-29.
- BOROJEVIĆ S. (1965): Način nasleđivanja i heritabilnost kvantitativnih svojstava u ukrštanjima raznih sorti pšenice. *Savremena polj.*, 7-8, 587-606.
- DUA P.R. and P.T. YADOVA (1985): Genetics of yield and its components in sunflower (*Helianthus annuus* L.). Proc. of 11th Int. Sunf. Conf., 627-632, Mar del Plata, Argentina
- EL-HITY M.A. (1992): Genetical analysis of some agronomic characters in sunflower (*Helianthus annuus* L.). Proc. of 13th Inter. Sunf. Conf., 1118-1128, Pisa, Italy
- ESENDAL E. and N. KANDEMIR (1996): Effects of row spacing on sunflower (*Helianthus annuus* L.) yield and other characteristics. Proc. of 14th Inter. Sunf. Conf., 369-374, 12-20, Beijing/Shenyang, China.
- FICK G.N. (1978): Breeding and genetics of sunflower. *Sunflower science and Technology, Agronomy* 19, 279-337.
- GIRARAJ K., S. VIDYASHANKART, M.N. VENKATARAM, and S. SEETHARAM (1979): Path coefficient analysis of seed yield in sunflower. *The Sunflower Newsletter*, 3, 10-12.

- GRIFFING B.A. (1956): A generalised treatment of the use of diallel cross in quantitative inheritance. *Heredity*, 10, 31-50.
- HADŽIVUKOVIĆ S. (1991): Statistički metodi. Drugo prošireno izdanje. Poljoprivredni fakultet, Novi Sad.
- IVANOV P. and Y. STOYANOVA (1980): Studies on the genotypic and phenotypic variability and some correlations in sunflower (*Helianthus annuus* L.). Proc of 9th Inter. Sunfl. Conf., 336-342, Tooremolinos, Espana
- JOKSIMOVIĆ J., J. ATLAGIĆ i D. ŠKORIĆ (2000): Efekat gena i kombinacione sposobnosti za prečnik glave kod nekih inbred linija suncokreta. Selekcija i semenarstvo, 1-2, 45-49
- KOVAČIK A. and V. ŠKALOUD (1971): Sunflower genetics and its application in sunflower selection. *Genetica a šlechtění*, 7, 29-54, Praha, Czechoslovakia
- KOVAČIK A. and V. ŠKALOUD (1972): Combining ability and prediction of heterosis in sunflower (*Helianthus annuus* L.). *Sci. Agriculture Bohemoslovaca*, 4 (4), 263-273
- KUMAR A.A., M. GANESH, and P. JANILA (1998): Combining ability analysis for yield and yield contributing characters in sunflower (*Helianthus annuus* L.). *Annals of Agr. Research*, 19, 437-440.
- MARINKOVIĆ R. (1984): Način nasleđivanja prinosa semena i nekih komponenti prinosa ukrštanjem raznih inbred linija. Doktorska disertacija, Univerzitet u Novom Sadu, Poljoprivredni fakultet.
- MARINKOVIĆ R. (1987): Analiza komponenti prinosa semena suncokreta (*Helianthus annuus* L.) koeficientom putanje I. Abstrakt, III Kongres genetičara Jugoslavije sa međunarodnim učešćem, Ljubljana, Jugoslavija.
- MARINKOVIĆ R. (1992): Path-coefficient analysis of some yield components of sunflower (*Helianthus annuus* L.). *Euphytica*, 60, 201-205.
- MARINKOVIĆ R. (1993): Combining ability of some inbred sunflower (*Helianthus annuus* L.) lines. *Indian J. Genet.*, 53, 299-304.
- MARINKOVIĆ R., B. DOZET i D. VASIĆ (2003): Oplemenjivanje suncokreta (Monografija), Školska knjiga, Novi sad, 368 str.
- MARINKOVIĆ R. i D. ŠKORIĆ (1990): Nasleđivanje prečnika glave i broja cvetova po glavi u ukrštanjima raznih inbred linija suncokreta (*Helianthus annuus* L.). *Uljarstvo*, 27 (1-2), 22-27.
- MATHER K. and J.L. JINKS (1982): Biometrical genetics. Third Edton, Chapman and Hall, London, England, 65-83.
- MICHELE R., R. Vincenzo, and F. FRANCESCO (1996): Effects of sowing on sunflower yield, oil and plant characteristics in Southern Italy. Proc. of 14th Inter. Sunf. Conf., 345-350, Beijing/Shenyang, China.
- MOROZOV V.K. (1947): Selekcija podsolnečnika v SSSR. (knjiga), Pišćepromizdat, Moskva
- PETAKOV D. (1994): Correlation and heritability of some quantitative characters in sunflower diallel crosses. Symposium on breeding of oil and protein crops, 162-164, Albena, Bulgaria
- PUNNIA M.S. and H.S. GILL (1995): Correlation and path coefficient analysis for seed yield traits in sunflower (*Helianthus annuus* L.). *Plant. Breed. Abs.*, 65, 915.
- PUTT ED (1965): Heterosis, combining ability and predicted synthetics from a diallel cross in sunflower (*Helianthus annuus* L.). *Can. J. Pl. Sci.*, 46, 59-67
- RAO N.M. and H.S. SINGH (1977): Inheritance of some quantitative characters in sunflower (*Helianthus annuus* L.). *Pak. J. of Rfs.* 2, 144-146.
- SCHUSTER W. (1964): Inzucht und Heterosis bei der Sonnenblume (*Helianthus annuus* L.). Willhem Schmitz-Verlag, 1-135, Giessen
- ŠEĆEROV-FIŠER V., J. Atlagić, and R. MARINKOVIĆ (1992): Inheritance of the head diameter in ornamental sunflower. Proc. of 13th Int. Sunf. Conf., 372-378. Pisa, Italy
- SETTY K.L.T. and B. SINGH (1977): Line x tester analysis of combining ability in sunflower. *Pak. J. Rfs.* 1, 23-26.
- SINDAGI S.S., R.S. KULKARNI, and A. SEETHARAM (1979): Line x tester analysis of the combining ability in sunflower (*Helianthus annuus* L.). *The Sunfl. Newsletter*, 3, 11-12.

- ŠKORIĆ D., T. VREBALOV, T. ČUPINA, J. TURKULOV, R. MARINKOVIĆ, S. MAŠIREVIĆ, J. ATLAGIĆ, L. TADIĆ, R. SEKULIĆ, D. STANOJEVIĆ, M. KOVAČEVIĆ, V. JANČIĆ i Z. SAKAČ (1989): Suncokret (monografija), Nolit, 613str.
- ŠKORIĆ D. (1975): Mogućnost korišćenja heterozisa na bazi muške sterilnosti kod suncokreta. Doktorska disertacija, Univerzitet u Novom Sadu, Poljoprivredni fakultet.
- ŠKORIĆ D., M. MARINKOVIĆ, S. JOCIĆ, D. JOVANOVIĆ i N. HLADNI (2002): Dostignuća i dalji pravci u oplemenjivanju suncokreta i izbor hibrida za setvu u 2002 godini, Zbornik radova Naučnog instituta za ratarsvo i povrtarstvo, Sveska 36, 147-160.
- STOJANOVA J., P. IVANOV, and J. GEORGIEV (1971): Nasledovanja na njakoj priznaci v F₁ pri slančogleda. Genetika i selekcija, 1, 3-14, Sofia.
- SUZER S. and I. ATAKISI (1993): Yield components of sunflower hybrids of different height. Helia, 16., 35-40.
- VOLF V.G. and L.P. DUMANČEVA (1973): Pojavlenije geterozisa u gibridov pervogo pokolonenija podsolnečnika. Geterozis kulturnih rastenij. Rez., 40. Varna

**KOMPONENTE FENOTIPSE VARIJABILNOSTI ZA PREČNIK GLAVE
SUNCOKRETA**Nada HLADNI, Dragan ŠKORIĆ i Marija KRALJEVIĆ-BALALIĆ¹

Naučni institut za ratarstvo i povrtarstvo, 21000 Novi Sad

¹ Poljoprivredni fakultet, 21000 Novi Sad, Srbija i Crna Gora

Izvod

Suncokret je u mnogim zemljama u svetu i kod nas osnovna biljka za proizvodnju jestivog ulja. Visina biljke, veličina, forma i položaj glave na stablu, broj listova, njihova veličina, trajanje i raspored na biljci imaju važnu ulogu u definisanju optimalne arhitekture hibrida suncokreta (ŠKORIĆ, 1975; 1989; 2002). U cilju praćenja načina nasleđivanja i efekta gena prečnika glave suncokreta u F₁ i F₂ generaciji izvršena su dialelna ukrštanja isključujući recipročna sa šest genetski divergentnim inbred linijama suncokreta. Dobijene su značajne razlike u srednjim vrednostima za ispitivano svojstvo. Prečnik glave se u F₁ generaciji nasleđivao superdominantno jedino se u četiri ukrštanja javila dominacija, dok se u F₂ generaciji nasleđivao dominantno, a superdominacija se zadržala u četiri kombinacije. Glavni deo genetske varijanse čini dominantna komponenta, a način nasleđivanja prečnika glave je superdominacija uzevši u obzir sve kombinacije ukrštanja u obe generacije (F₁ i F₂). Ova istraživanja mogu biti od značaja u stvaranju visoko prinostnih genotipova suncokreta.

Primljeno 4. XI 2003.
Odobreno 29. XII 2003.