

**GENOTYPE SPECIFICITY IN NITROGEN NUTRITION
OF MALTING BARLEY**

Vera POPOVIĆ¹, Đorđe GLAMOČLIJA², Miroslav MALEŠEVIĆ¹,
Jela IKANOVIĆ², Gordana DRAŽIĆ³, Marija SPASIĆ⁴ and Saša STANKOVIĆ⁵

¹Institute of Field and Vegetable Crops, Novi Sad, Serbia

²Faculty of Agriculture, Belgrade, Serbia

³Institute of Applied Ecology, „Futura“, Belgrade, Serbia

⁴Agromaster Company, Kragujevac, Serbia

⁵Technological Research Center, Zaječar, Serbia

Popović V, Đ. Glamočlija, M. Malešević, J. Ikanović, G. Dražić, M. Spasić and S. Stanković (2011): *Genotype specificity in nitrogen nutrition of malting barley*. - Genetika, Vol 43, No. 1, 197 -204.

A three-year trial (2003-2005) was conducted under agro ecological conditions of Timočka Krajina (the experiment farm of Technological Research Center in Zaječar). Research object were six malting barley genotypes, which were top-dressed with the following amounts of nitrogen in the course of growing season: 40, 60, 80 and 100 kg ha⁻¹. A non-fertilized variant served as a control. The obtained results indicated that the tested genotypes reacted to increased amounts of nitrogen by changing their morphological and biological characteristics as well as

Corresponding author: Vera Popović, Institute of Field and Vegetable Crops, Novi Sad, Tel: +381 21 4898481; E-mail: vera.popovic@ifvcns.ns.ac.rs

the technological values of grain. The effectiveness of the applied nitrogen depended significantly on the distribution of rainfall in periods of highest water uptake by malting barley.

Key words: genotype, grain quality, malting barley, morphological and biological characteristics, nitrogen fertilization

INTRODUCTION

The economic importance of malting barley is due to the use of barley grain in the industries of beer and hard liquors. Quality barley for malting industry should contain more than 60% of starch and less than 12% of total proteins. Bulk weight should be over 65 kg, and 1000-grain weight above 40 g. Modern barley breeding is aimed at the development of genotypes that meet the requirements of processing industry (DJURIĆ *et al.*, 2009). In addition to genotype, has a major impact and properly balanced nutrition, adapted to actual soil fertility and barley requirement for nitrogen, determine the quality of grain (GLAMOČLIJA *et al.*, 1998). The effect of nitrogen on barley characteristics depends on N amount applied and weather conditions in the growing year. Increased amounts of nitrogen curtail its positive effects on spike length, number of grains per spike, 1000-grain weight and yield level. However, the protein content in grain keeps increasing with nitrogen dose, which lowers the quality of malting barley (Malešević *et al.*, 2010). Conversely, PRŽULJ and MOMČILOVIĆ (2001) claimed that intensive nitrogen fertilization significantly increases crop yield, while grain quality and its utility depend on the genotype. The rate of utilization of applied nitrogen depends also on environmental conditions. In years with less rainfall in the spring, plants better utilize the top-dressed nitrogen (PAGOLA *et al.*, 2009). Many authors (KENDALL, 1996; PRŽULJ *et al.*, 1998; GLAMOČLIJA *et al.*, 1999; PAUNOVIĆ *et al.*, 2006, 2008; MADIĆ *et al.*, 2009) reported that an increased number of spikes increases the yield of malting barley. The chemical composition of vegetative biomass depends on the variety and intensity of plant nutrition, but also from dynamics of individual phenological phases (UZUN, 2010).

GRANT *et al.* (1991) claimed that, on soils with low N supplies, malting barley responds well to N fertilizer, exhibiting increases in yield and protein content. However, too much nitrogen can increase protein beyond levels acceptable to the malting industry's standards. New barley genotypes have a high genetic yield potential, getting close to wheat in productivity. Therefore, the barley is considered a prospective crop for Serbia, especially from the point of the global climate change which has not avoided the agricultural regions of Serbia. It is obvious that the impact of drought on the agricultural crops keeps intensifying. Therefore, it is recommended to rely on small grains in the agricultural production because they are tolerant to drought. Barley is highly tolerant to drought and high temperature and it belongs to the crops capable of mitigating the harmful effects of the climate change.

The objectives of this study were to find the most suitable assortment of barley cultivars and determine their genetic yield potential when grown under the agroecological conditions of eastern Serbia.

MATERIALS AND METHODS

The three-year trial (2003-2005) was carried out in the agroecological conditions of Timočka Krajina (the experiment farm of Technological Research Center in Zaječar), on the limeless smonitza soil. Agrochemical analyses showed that the chemical reaction of the soil was slightly acidic (pH - 5.80), and that the soil was provided with phosphorus (P_2O_5) and potassium (K_2O) (17.5% and 29.95%, respectively), medium provided the nitrogen (N) (0.12%), and rich in humus (3.08%).

The research object were genotypes of winter malting barley *Kristal* (G_1), *Premijum* (G_2), *NS -519* (G_3), *NS-525* (G_4), and two lines, *ZA-82/1* (G_5) and *ZA -12/1* (G_6). The experimental plants received the following amounts of mineral fertilizers. During seedbed preparation, 30 kg ha⁻¹ of N, 80 kg ha⁻¹ of P_2O_5 and 60 kg ha⁻¹ of K_2O were applied. Top dressing was performed in the course of the winter season, applying 40, 60, 80 and 100 kg ha⁻¹ of N (variants N_1 , N_2 , N_3 and N_4 , respectively). The control variant was not fertilized (N_0).

Phenological stages were observed in the course of the growing season and plant samples for analyses were taken at the stage of full maturity. The following parameters were measured: plant height, number of spikes per m², spike length, 1000-grain weight and hectolitre weight. All variants were sampled for determination of germinability and germination energy.

The obtained data were processed by the analysis of variance.

The annual amounts of rainfall in the three years (638 mm, 665 mm and 672 mm) were above the long-term average (586 mm). The distribution of rainfall in the first year of study was unfavourable. Long periods of drought at the stages of tillering, shooting and heading significantly affected the development of plants. The highest rainfall during growing season occurred in the second year. Most favourable distribution of rainfall was in the third year. The temperature regime was within the long-term average values for the studied region.

RESULTS AND DISCUSSION

Barley stem height was affected by increased nitrogen and genotype in all three years of study. The effect of the treatments depended on the weather conditions during the barley growing season (Table 1). PECIO and BICHONSKI (2002) emphasize the high effect of nitrogen on the total growth of barley plants. TAMM and TAMM (2000) reported significant correlation between varieties and environmental conditions.

The number of spikes per meter square, or intensity of tillering, was lowest in the first year (318). In the third year, the tillering was 2 times higher, and in the second year 2.5 times higher than in the first year. Nitrogen fertilizers affected the intensity of tillering in all three years, with differences among genotypes being significant only in the second year (Table 2). VEIGH and RAJKAI (2006) emphasize an optimal nitrogen rate to be applied in barley nutrition should be identified.

Table 1. Stem height, cm

Genotype	2002/3						2003/4						2004/5						
	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	\bar{X}
G ₁	33	38	41	44	46	40.4	66	77	82	85	83	78.6	57	64	69	72	73	67.0	62.0
G ₂	32	38	41	44	46	40.2	67	73	77	78	80	75.0	58	62	66	69	70	65.0	60.1
G ₃	32	42	44	45	45	41.6	68	76	79	81	81	77.0	59	64	66	70	70	65.9	61.5
G ₄	34	43	44	46	46	42.6	68	76	80	83	84	78.2	57	65	70	74	75	68.2	63.0
G ₅	33	37	41	42	44	39.4	64	76	77	79	80	75.2	56	61	66	67	68	63.6	59.4
G ₆	35	41	44	45	46	42.2	67	76	84	83	84	78.8	57	65	71	75	76	68.8	62.3
Average	33.2	39.8	42.5	44.3	45.5	41.1	66.7	75.7	79.8	81.5	82.0	77.1	57.3	63.5	68.0	70.5	72.0	66.3	61.5

Characteristic tested	Test	2002/3			2003/4			2004/5		
		Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB
Stem height	F test	**	**	**	**	**	**	NS	**	*
	LSD 5%	1.1603	0.8560	2.2438	3.6093	1.6044	5.1384	1.4038	1.1094	2.8532
	1%	1.6503	1.1268	3.0493	5.1337	2.1178	7.1178	1.9967	1.4603	3.8677

Table 2. Spike number per m²

Genotype	2002/3						2003/4						2004/5						
	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	\bar{X}
G ₁	253	283	345	347	349	315	652	787	847	861	860	801	560	655	679	699	695	658	591
G ₂	243	285	349	352	352	316	652	833	860	880	881	821	568	656	685	695	697	660	599
G ₃	248	281	348	352	352	316	633	795	860	878	884	810	555	663	688	700	700	661	596
G ₄	246	295	351	352	355	320	652	790	855	872	881	810	557	651	689	695	701	659	596
G ₅	257	289	352	360	357	323	648	801	855	887	889	815	546	652	691	701	696	657	598
G ₆	253	288	348	355	353	319	662	779	845	864	873	805	550	660	688	696	702	659	594
Average	250	287	349	353	353	318	649	797	853	873	878	810	556	656	686	697	698	659	596

Characteristic tested	Test	2002/3			2003/4			2004/5		
		Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB
Spike number	F test	NS	**	NS	**	**	*	NS	**	NS
	LSD 5%	12.0048	9.4283	24.2884	10.6032	8.7004	22.3140	10.9456	9.3269	23.5391
	1%	17.0750	12.4107	32.9314	15.0815	11.5578	30.1967	15.5685	12.2772	31.8224

Spike length was found to depend in all three years on the genotype and the amount of nitrogen, while the interaction of these two factors was significant only in the third year (Table 3).

Barley is a kind of small grains with the highest tillering coefficient. It depends on the genotype, nitrogen nutrition of plants, as well as on environmental conditions (MADIĆ *et al.*, 2006). On the other hand, VEIGH and RAIKAI (2006) claimed that optimum soil moisture in the initial stages of growth significantly affects the development of roots. Well-rooted plants are more intense tillering of the increased nitrogen supply.

The weight of 1000 grains is an important indicator of the utility of malting barley. Its value was lowest in the first year and the overall average was below 40 g. In the second and third years, the 1000'grain weight was above 40 g in all six genotypes, which meant that we obtained quality raw materials for the beer industry. Variations of the treatment groups were highly significant per genotype and per nitrogen nutrition of plants (Table 4). According to PRŽULJ and MOMČILOVIĆ (2002)

and PRŽULJ *et al.* (1998) the optimal amount of nitrogen for malting barley nutrition, from the standpoint of grain size, is 80 kg ha⁻¹.

Table 3. Spike length, cm

Genotype	2002/3						2003/4						2004/5						
	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	\bar{X}
G ₁	7.1	7.3	8.0	8.3	8.5	7.9	8.7	9.6	10.5	10.6	10.7	10.0	7.7	8.5	8.6	9.1	9.5	8.7	8.9
G ₂	7.0	7.7	8.3	8.4	8.6	8.0	8.7	9.8	10.6	10.9	10.9	10.2	7.4	8.4	9.0	9.4	9.4	8.8	9.0
G ₃	7.2	8.1	8.7	8.7	9.0	8.3	8.7	10.2	10.8	11.4	11.2	10.5	7.8	9.2	9.8	10.2	10.5	9.5	9.4
G ₄	7.1	7.8	8.4	8.8	8.9	8.2	8.3	9.8	10.5	10.9	10.1	7.7	8.6	9.2	9.4	9.4	8.9	9.1	
G ₅	7.0	7.6	8.6	8.8	8.9	8.2	8.5	10.0	10.6	10.8	10.7	10.1	7.9	8.6	9.2	9.4	9.4	8.9	9.1
G ₆	7.1	8.0	8.9	9.3	9.4	8.5	9.3	10.2	11.3	11.7	11.9	10.9	8.0	9.2	10.0	10.4	10.3	9.6	9.7
Average	7.1	7.8	8.5	8.7	8.9	8.2	8.7	9.9	10.7	11.1	11.1	10.3	7.8	8.8	9.3	9.7	9.8	9.1	9.2

Characteristic tested	Test	2002/3			2003/4			2004/5		
		Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB
Spike length	F test	**	**	NS	*	**	NS	**	**	*
	LSD 5%	0.2619	0.1661	0.4568	0.4380	0.4121	1.0175	0.2330	0.1820	0.4695
	1%	0.3724	0.2186	0.6244	0.6229	0.5425	1.3709	0.3314	0.2395	0.6367

Table 4. 1000 - grain weight, g

Genotype	2002/3						2003/4						2004/5						
	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	\bar{X}
G ₁	36.4	37.7	39.0	39.2	39.9	38.5	40.0	43.5	44.7	45.9	46.0	44.0	41.1	44.5	46.2	46.7	47.1	45.1	42.5
G ₂	36.9	37.6	39.2	40.0	40.8	38.9	40.7	41.4	43.4	43.9	42.6	41.6	43.8	45.3	45.7	46.0	44.5	42.0	
G ₃	37.6	40.5	42.0	42.3	43.2	41.1	40.6	44.5	46.0	47.3	47.1	45.1	45.7	46.5	48.1	49.5	49.7	47.9	44.7
G ₄	36.4	42.2	42.5	43.2	43.6	41.6	40.0	44.8	47.7	47.1	48.2	45.6	41.1	47.2	49.3	50.7	51.8	48.0	45.1
G ₅	38.0	38.7	39.6	40.0	40.2	39.3	40.7	43.3	44.4	44.9	45.5	43.8	42.1	46.7	48.0	49.1	49.5	47.1	43.4
G ₆	37.3	38.1	38.6	39.1	39.4	38.5	41.2	42.3	43.6	44.1	44.4	43.1	41.2	44.0	46.2	46.4	46.7	44.9	42.2
Average	37.1	39.1	40.2	40.6	41.2	39.7	40.5	43.3	45.0	45.6	45.9	44.0	42.1	45.5	47.2	48.0	48.5	47.8	43.8

Characteristic tested	Test	2002/3			2003/4			2004/5		
		Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB
1000-grain weight	F test	**	**	**	**	**	**	**	**	*
	LSD 5%	0.3801	0.3095	0.7899	0.4558	0.5602	1.3223	0.9658	0.8543	2.1376
	1%	0.5406	0.4074	1.0696	0.6483	0.7374	1.7679	1.3737	1.1246	2.8861

Bulk weight is also an important indicator of technological value of grain. In the first year, it was near the lower threshold of the acceptable quality (the overall average value of 64.2 kg). In the second and third years, the grain was of excellent quality, the bulk weights being 69.4 and 72.1 kg, respectively. Highest fluctuations in the values of bulk weight were registered in the first year when significant differences were recorded between the genotypes, the amounts of nitrogen and the interaction of these two factors. There were no significant differences in the second year, while in the third year the genotype x nitrogen interaction was non-significant (Table 5).

The barley seed from the second year had the lowest germination energy (84.9%). The respective values in the third and first years were 89.2% and 92.3%. The three-year overall average value and the annual average values depended to a large measure on the genotype and amount of nitrogen. The interaction genotype x

nitrogen was non-significant only in the first year (Table 6). Significant effect of weather conditions on seed quality was also indicated in a study of GLAMOČLIJA *et al.* (1998). According to JACKSON (2000), environmental factors such, as drought stress, that occur late in the season can adversely affect grain yield, and, in particular, quality characteristics. Kernel plumpness characteristics are strongly related to yield potential and available nitrogen.

Table 5. Bulk weight, kg

Geno- type	2002/3						2003/4						2004/5						
	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	\bar{X}
G ₁	61.4	62.4	63.0	65.5	64.3	62.9	69.0	69.9	72.0	72.7	72.8	71.3	65.1	66.4	68.8	69.5	70.1	68.0	67.4
G ₂	62.5	64.5	65.8	67.2	76.5	65.5	68.4	71.1	73.9	74.6	75.1	72.6	66.5	67.9	68.9	70.7	71.6	69.1	69.1
G ₃	62.2	64.0	65.4	65.4	65.9	64.6	68.0	74.2	73.5	72.6	73.4	72.3	65.6	66.7	69.5	70.4	70.7	68.6	68.5
G ₄	62.1	64.3	65.9	66.9	67.4	65.3	68.2	70.1	73.5	73.9	74.7	72.0	65.8	69.4	72.6	73.6	74.1	71.1	69.5
G ₅	62.5	63.9	64.4	64.8	65.8	64.3	68.4	72.6	71.9	74.1	74.0	72.2	67.7	69.5	72.3	71.9	73.6	71.0	69.2
G ₆	54.1	61.2	61.8	62.8	63.6	62.7	68.2	73.3	72.9	73.2	73.8	72.3	66.4	67.5	68.2	69.7	70.4	68.4	67.8
Average	60.8	63.4	64.4	65.4	67.3	64.2	68.4	71.9	73.0	73.5	73.9	72.1	66.2	67.9	70.1	71.0	71.8	69.4	68.6

Characteristic tested	Test	2002/3			2003/4			2004/5		
		Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB
Bulk weight	F test	**	**	**	NS	NS	NS	**	**	NS
	LSD 5%	0.9102	0.6853	1.7860	1.6928	1.2172	3.2147	0.6484	0.8014	1.8902
	1%	1.2946	0.9021	2.4253	2.4077	1.6022	4.3730	0.9223	1.0550	2.5268

Table 6. Germination energy, %

Genotype	2002/3						2003/4						2004/5						
	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	N ₀	N ₁	N ₂	N ₃	N ₄	\bar{X}	\bar{X}
G ₁	88.7	90.0	93.7	94.7	95.7	92.5	77.0	78.3	82.0	84.0	85.0	81.3	85.7	87.7	90.0	92.3	94.0	89.9	87.9
G ₂	89.7	90.7	93.7	94.7	96.3	93.0	79.7	81.7	86.3	87.7	89.3	84.9	88.3	88.3	91.7	93.0	94.7	91.2	89.7
G ₃	90.3	90.0	90.7	92.7	92.7	91.3	78.7	76.0	78.3	81.3	84.0	79.7	85.3	84.0	87.0	89.0	89.3	86.9	86.0
G ₄	90.7	90.3	94.0	93.7	94.7	92.7	87.0	85.3	89.0	90.3	91.0	88.5	87.0	85.3	89.0	90.3	91.0	88.5	89.9
G ₅	89.0	90.7	93.7	94.7	96.0	92.8	86.7	88.7	90.0	91.3	92.0	89.7	86.7	88.3	90.0	91.3	92.0	89.7	90.7
G ₆	88.0	89.0	91.0	94.0	95.3	91.5	87.0	79.7	83.3	87.3	88.3	85.1	88.3	86.0	89.0	90.7	91.0	89.0	88.5
Average	89.4	90.1	92.8	94.1	95.1	92.3	82.7	81.6	84.8	87.0	82.3	84.9	86.9	86.6	89.5	91.1	92.0	89.2	88.8

Characteristic tested	Test	2002/3			2003/4			2004/5		
		Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB	Genotype	Nitrogen	AxB
Germination energy	F test	**	*	NS	**	**	**	**	**	**
	LSD 5%	0.9741	0.7622	1.9655	1.1300	0.9143	2.3371	0.8206	0.9149	2.1905
	1%	1.3856	1.0033	2.6653	1.6072	1.2035	3.1653	1.1672	1.2043	2.9364

CONCLUSION

The three-year research results of genotypic specificity in nitrogen nutrition of malting barley allow us to draw the following conclusions.

- In the year with the unfavourable distribution of rainfall, nitrogen top dressing increased the average stem height by 29.1%. In the favourable years, the increases were 19.5% and 19.6%.

- The overall average number of spikes per unit area was highest when 100 kg/ha of nitrogen were applied (643). The highest effect was registered in the first, dry year.

- Water regime affected the length of the spike. Under dry conditions and on the overall average, the genotype NS-519 and the line 12 // formed a significantly longer spike.

- The genotypes *NS-519* and *NS-525* had largest grains whose bulk weight exceeded 40 grams even in the dry year.
- Bulk weight of grain was very good on the whole; as it was less dependent on the distribution of rainfall. In the year with the most favourable water regime, the studied factors did not influence this parameter. The line *82/1* was distinguished throughout the study period for highest bulk weight.
- Abundant RAINFALL during the barley maturation in the second year adversely affected the germination energy, which was below the standard value required by the beer industry.

ACKNOWLEDGMENTS

Support by the Ministry of Science and Technological Development, Republic of Serbia, to this study (project: TR 20069) is gratefully acknowledged.

Received, December 15th 2010

Accepted, March 14th 2011

REFERENCES

- DURIĆ, N., V., TRKULJA, S., PRODANOVIĆ (2009). Breeding and production of malting barley created at the PKB Institute. Proceedings of the Institute PKB Agroekonomik, No 1-2, pp 21-26.
- GLAMOČLIJA, Đ., J., LAZAREVIĆ, D., KOVAČEVIĆ and L., RUŽIČIĆ (1998). Effects of nitrogen top dressing and microelements foliar fertilization on the yield and yield components of malting barley. Proceedings, Breeding of Small Grains, Kragujevac, 401-405.
- GLAMOČLIJA, Đ., J. LAZAREVIĆ, LJ. PRIJIĆ, D. KOVAČEVIĆ, D. ŽIVKOVIĆ (1999). Influence of nitrogen nutrition and micronutrients on some morphological characteristics of malting barley. Journal of Scientific Agricultural Research, Vol. 60, No 210, pp. 25-33, Beograd.
- GRANT, C. A., L. E. GAUER, D. T. GEHL, and L. D. BAILEY (1991). Yield response of semidwarf and conventional height barley cultivars to nitrogen fertilization under varying moisture conditions. Canadian Journal Plant Sciences, No 71, pp. 361-371.
- JACKSON, G. D. (2000). Nitrogen Fertilization of Dryland Malt Barley for Yield and Quality. Fertilizers Facts, Montana State University
- KENDALL, N. T. (1996). North American brewing industry requirements. y: Proceedings of V International oat Conference and VII International Berley Genetics Symposium, Saskatoon: University of Saskatchewan, 26-31
- MADIĆ, M., D. KNEŽEVIĆ, A. PAUNOVIĆ and N. BOKAN (2006). Variability and inheritance of tillering in barley hybrids, Genetika, Vol. 38, No. 3, pp. 193-202.
- MADIĆ, M., A. PAUNOVIĆ, D. KNEŽEVIĆ and V. ZEČEVIĆ (2009): Grain Yield And Yield Components Of Two-Row Winter Barley Cultivars And Lines, Acta agriculturae Serbica, Vol. 14, 27, 17-22.
- MALEŠEVIĆ, M., Đ., GLAMOČLIJA, N., PRŽULJ, V., POPOVIĆ, S., STANKOVIĆ, T. ŽIVANOVIĆ, and A., TAPANAROVA (2010). Production characteristics of different malting barley genotypes in intensive nitrogen fertilization. Genetika, Vol.42, No. 2, pp. 323-330.
- PAGOLA, M., O. RUBCIN, I. IRIGOYEN, H. BUSTINCE, E. P. BARRENECHEA, C. APARICIO-TEJO, L. LAMSFUS BERTA (2009). New method to assess barley nitrogen nutrition status based on image colour analysis: Comparison with SPAD-502. Computers and Electronics in Agriculture, Vol. 65, Issue 2, Pp. 213-218.
- PAUNOVIĆ, A., D. KNEŽEVIĆ, and M. MADIĆ (2006): Genotype variations in grain yield of spring barley depending on sowing density, Genetika, Vol. 38 No. 2., 107-114.
- PAUNOVIĆ, A., M. MADIĆ, D. KNEŽEVIĆ and M. BIBERDŽIĆ (2008). Nitrogen and seed density effects on spike length and grain weight per spike in barley. Cereal Research Communications, Vol. 36, 75-78.
- PECIO, A. and A. BICHONSKI (2002). Productive results of malting barley nitrogen fertilization. Pamietnik-Pulawski, No.130 (2), 557-564.

- PRŽULJ, N. and V. MOMČILOVIĆ (2002). NS barley varieties for the agroecological conditions of Southeastern Europe. *Proceedings of the Institute of Field and Vegetable Crops, Novi Sad, No 36*, pp 271-282.
- PRŽULJ, N., S. DRAGOVIĆ, M. MALEŠEVIĆ, V. MOMČILOVIĆ and N. MLADENOV (1998). Comparative performanse of winter and spring malting barleys in semiarid growing conditions. *Euphytica 101*: 377-382.
- TAMM, I. and Ü. TAMM (2002). Genetic and environmental variation of malting barley and oat grain quality characteristics. *Agriculture, 78*, pp. 51 - 57.
- UZUN, F. (2010). Changes in hay yield and quality of bulbous barley at different phenological stages. *Turkish Journal of Agriculture and Forestry, Vol. 34*, pp. 1-9.
- VEIGH, K. R. and K. RAIKAI (2006). Rooth growts and nitrogen use efficiency of spring barley in drying soil. *Cereal Research Communications, 34* (1), pp. 267-270.

GENOTIPSKE SPECIFIČNOSTI PIVARSKOG JEČMA U USLOVIMA INTENZIVNE ISHRANE AZOTOM

Vera, POPOVIĆ¹, Đorđe GLAMOČLIJA², Miroslav, MALEŠEVIĆ¹,
Jela IKANOVIĆ², Gordana DRAŽIĆ³, Marija SPASIĆ⁴ i Saša STANKOVIĆ⁵

¹ Institut za ratarstvo i povrtarstvo, Novi Sad, Srbija

² Poljoprivredni fakultet, Beograd, Srbija

³ Institut za primenjenu ekologiju, „Futura“, Beograd, Srbija

⁴ Agromaster Kompanija, Kragujevac, Srbija

⁵ Centar za poljoprivredna i tehnološka istraživanja, Zaječar, Srbija

I z v o d

Trogodišnji ogledi (2003-2005) su izvedeni u Centru za poljoprivredna i tehnološka istraživanja u Zaječaru. Materijal istraživanja bilo je šest genotipova pivarskog ječma koji su tokom vegetacionog perioda prihranjivani sledećim količinama azota 40, 60, 80 i 100 kg ha⁻¹. Kao kontrola poslužila je varijanta bez prihranjivanja. Dobijeni rezultati pokazali su da genotipovi reaguju na povećane količine azota promenom morfoloških i bioloških osobina, kao i promenama tehnološke vrednosti semena. Efekti upotrebljenog azota značajno zavise od rasporeda padavina u periodima najveće potrošnje vode.

Primljeno 15. XII. 2010.

Odobreno 14. III. 2011.