

## MOLECULAR TOOLS FOR UTILIZATION OF MITOCHONDRIAL DIVERSITY IN FABA BEAN (*Vicia faba*)

Jelena M. ALEKSIĆ<sup>1</sup>, Jovanka MILJUŠ-ĐUKIĆ<sup>1</sup>, Živko JOVANOVIĆ<sup>1</sup>, Branko TOMIĆ<sup>2</sup>  
and Bojana BANOVIĆ<sup>1</sup>

<sup>1</sup> - Laboratory for Plant Molecular Biology, Institute of Molecular Genetics and Genetic Engineering (IMGGE), University of Belgrade, Belgrade, Serbia

<sup>2</sup> - Laboratory for Molecular Biology, Institute of Molecular Genetics and Genetic Engineering (IMGGE), University of Belgrade, Belgrade, Serbia

Aleksić M. J., J. Miljuš-Đukić, Ž. Jovanović, B. Tomić and B. Banović (2014): *Molecular tools for utilization of mitochondrial diversity in faba bean (Vicia faba)*. - Genetika, Vol 46, No. 3, 745- 762.

We performed *in silico* PCR analyses utilizing complete mitochondrial (mtDNA) genome sequences of faba bean (*Vicia faba*) and two related species, *Vigna angularis* and *Vigna radiata*, currently available in GenBank, to infer whether 15 published universal primer pairs for amplification of all 14 *cis*-spliced introns in genes of NADH subunits (*nad* genes) are suitable for *V. faba* and related species. Then, we tested via PCR reactions whether seven out of 15 primer pairs would generate PCR products suitable for further manipulation in 16 genotypes of *V. faba* representing all botanical varieties of this species (*major*, *minor*, *equina* and subsp. *paucijuga*) of various levels of improvement (traditional and improved cultivars) originating from Europe, Africa, Asia and south America. We provide new PCR primers for amplification of *nad1* intron 2/3 in *V. faba*, and demonstrate intraspecific variability in primary nucleotide sequences at this locus. Based on outcomes of both *in silico* predictions and PCR amplification, we report a set of PCR primers for amplification of five introns in *nad* genes that are promising molecular tools for future phylogeographic and other studies in this species for which unambiguous data on wild ancestors, centre of origin and domestication are lacking.

**Key words:** mitochondrial genome, NADH dehydrogenase, introns, *in silico* PCR, universal primers, faba bean

### INTRODUCTION

Faba bean (*Vicia faba* L.) is a grain crop legume used mainly for food and feed predominantly in eastern and northern Africa and western Asia (AKIBODE and MAREDIA, 2011).

---

**Corresponding author:** Jelena M. Aleksić, Laboratory for Plant Molecular Biology, Institute of Molecular Genetics and Genetic Engineering (IMGGE), University of Belgrade, Vojvode Stepe 444a, PO box 23, 11010 Belgrade, Serbia, E-mail: [aleksic\\_jelena@yahoo.com.au](mailto:aleksic_jelena@yahoo.com.au), [jelena.aleksic@imgge.bg.ac.rs](mailto:jelena.aleksic@imgge.bg.ac.rs)

Archaeological evidence reveals that *V. faba* is one of the first domesticated crops (TANNO and WILLCOX, 2006; see also DUC *et al.*, 2010 and references therein). However, its origin and wild ancestors are still unknown (MURATOVA, 1931; CUBERO, 1974; MAXTED *et al.*, 1991; MAXTED, 1995) and furthermore, its taxonomic placement and evolutionary relations are still unresolved despite numerous traditional morphology-based taxonomies and systematics (more than 20 since Linnaeus, reviewed by MAXTED, 1993) and available molecular phylogenies (e.g. POTOKINA *et al.*, 1999; VAN DE VEN *et al.*, 1993; FENNELL *et al.*, 1998; SCHAEFER *et al.*, 2012). New insights into these essential aspects of *V. faba* history, which are also required for breeding programs and crop improvement, may be gained via new approaches such as phylogeography (AVISE *et al.*, 1987; AVISE, 2009). Phylogeography is a relatively new discipline which makes inferences based on spatial arrangements of genealogically linked lineages detected mainly at haploid, non-recombining and predominantly uniparentally inherited cytoplasmic genomes (chloroplast - cpDNA and mitochondrial - mtDNA genomes in plants).

Given the lack of cpDNA diversity in *V. faba* (SHIRAN and MASHAYEKH, 2004; HAIDER *et al.*, 2012) and high mtDNA diversity obtained via PCR-RFLP method and random mtDNA clones used as probes for hybridisation (SCALLAN and HARMEY, 1996) as well as nucleotide and length variability at intron 2/3 of the first gene of the NADH dehydrogenase (*nad1* intron 2/3) in the genus *Vicia* (RYZHOVA *et al.*, 2012), it appears that mtDNA may be a promising tool not only for phylogeographic studies in *V. faba* that may shed more light on species origin and domestication but also for phylogenetic surveys that may resolve species evolutionary unfolding. However, utilization of maternally inherited (MOGENSEN, 1996) and slowly evolving mtDNA genome (WOLFE *et al.*, 1987; LAROCHE *et al.*, 1997) in diverse studies in *V. faba* may not be straightforward because plant mtDNA genomes are exceptionally variable in size (200 to 2.900 kb) and organization frequently altered by insertions, deletions and structural re-arrangements (LAROCHE *et al.*, 1997; see also NEGRUK, 2013 and references therein). This implies that it may be challenging not only to delineate informative regions but also to generate PCR products which can be used for further manipulation. This is because so-called universal primers, which are expected to generate PCR products in a vast majority of plant taxa (TABERLET *et al.*, 1991), may simply fail to amplify desired mtDNA regions in taxa of interest due to alternative organization of their genomes. Nonetheless, a recent report on complete mtDNA sequence of *V. faba* (NEGRUK, 2013) represents an essential milestone which would facilitate broader utilization of mtDNA variability in diverse studies in this species.

The aim of this study is to provide molecular tools for broader utilization of mtDNA diversity in diverse studies in *V. faba*. We firstly test via *in silico* PCR predictions whether universal mtDNA primers reported by DEMESURE *et al.* (1995), DUMOLIN-LAPEGUE *et al.* (1997) and DUMINIL *et al.* (2002) are suitable also for *V. faba* and related species *Vigna angularis* and *Vigna radita* whose complete mtDNA genome sequences are currently available in GenBank. We focus primarily on 15 primer pairs which amplify 14 *cis*-spliced introns in genes of NADH subunits (*nad* genes) because mtDNA non-coding regions are evolving faster than their adjoining coding regions, and introns in *nad* genes are amongst the most commonly used non-coding mtDNA regions in diverse studies in higher plants (e.g. FREUDENSTEIN and CHASE, 2001; WON and RENNER, 2003; ZHANG *et al.*, 2006; TOLLEFSRUD *et al.*, 2009; ALEKSIĆ and GEBUREK, 2010; HAVANANDA *et al.*, 2010; FROELICHER *et al.*, 2011; RYZHOVA *et al.*, 2012; LOCKWOOD *et al.*, 2013). Secondly, we test via PCR reaction whether seven out of 15 tested primer pairs would generate PCR products suitable for further manipulation in 16 genotypes of *V. faba* originating

from Europe, Africa, Asia and south America which represent all botanical varieties of this species (*major*, *minor*, *equina* and subsp. *paucijuga*) of various levels of improvement (traditional cultivars and improved cultivars). We provide new PCR primers for amplification of *nad1* intron 2/3 in *V. faba*, demonstrate intraspecific variability in primary nucleotide sequences at this locus, and based on outcomes of both *in silico* predictions and PCR amplifications, report a set of PCR primers for amplification of five introns in *nad* genes that are promising molecular tools for future phylogeographic and other studies in *V. faba*.

## MATERIALS AND METHODS

### *Complete mtDNA genome sequences and in silico PCR*

Mitochondrial genome sequences of *Vicia faba* cultivar Broad Windsor (GenBank accession number KC189947, NEGRUK, 2013), *Vigna radiata* (HM367685, ALVERSON *et al.*, 2011) and *Vigna angularis* (AP012599, NAITO *et al.*, 2013) have been downloaded from NCBI database on 14 January 2014. They were used for *in silico* PCR amplification of 14 cis-spliced introns in genes of NADH subunits (*nad1* introns 2/3 and 4/5; *nad2* introns 1/2, 3/4, and 4/5; *nad4* introns 1/2, 2/3 and 3/4; *nad5* introns a/b and d/e; and *nad7* introns 1/2, 2/3, 3/4 and 4/5) utilizing 15 universal primer pairs published by DEMESURE *et al.* (1995), DUMOLIN-LAPEGUE *et al.* (1997) and DUMINIL *et al.* (2002). *In silico* PCR was performed utilizing FastPCR 6.3 trial version (KALENDAR *et al.*, 2009). The maximal length of expected PCR products was set to 8 kb and allowed mismatch at 3'-end was set to two nucleotides. The positions of all primer binding sites of forward (F) and reverse (R) primers, the complementarity of primers and template DNA expressed through the percentage of binding nucleotides, melting temperatures (*T<sub>m</sub>*) of individual primers (in °C), annealing temperatures (*T<sub>a</sub>*) of primer pairs (in °C), and lengths of obtained PCR products were recorded in all legumes.

### *Plant material for PCR reactions*

Plant material used for PCR amplification of seven introns in *nad* genes comprised 16 genotypes of *V. faba* representing different botanical varieties (*major* – abbreviation Ma, *minor* – Mi, and *equina* - Eq) and subsp. *paucijuga* (Pau) which are traditional cultivars (TC) or improved cultivars (IC) originating from China (cultivar name: Kingpi – characteristics: Eq, TC), India (CH182 –Pau, TC), Ethiopia (E14b – Eq, TC), Morocco (BPL4225 – not available), Sudan (Silian – Mi, TC), Greece (Apsalos –Eq, IL), Italy (Violetta di policoro – Ma, TC), Spain (Fève Seville – Ma, TC), Denmark (Troy – Eq, IC), Finland (Mikko – Mi, TC), Germany (Diana – Mi, IC), United Kingdom (CH170 – Eq, IL), Peru (Rojo – Ma, TC) and Serbia (three traditional cultivars of *major* type - Bački Petrovac 2 and Deligrad, and of *minor* type – 159B). Seeds of all accessions, obtained from the French National Institute for Agricultural Research (INRA) *ex situ* collection, were sawn and grown to seedlings in a greenhouse at Institute of Molecular Genetics and Genetic Engineering (IMGGE). Leaves from young seedlings were collected and desiccated with silica gel for c. two weeks prior to DNA extraction. Total genomic DNA was extracted using CTAB method, modification according to ALEKSIĆ *et al.* (2012).

### *PCR primers and PCR amplification of seven introns in nad genes in V. faba*

Introns in *nad* genes (*nad1* intron 2/3; *nad4* introns 1/2 and 2/3; *nad5* intron d/e; and *nad7* introns 1/2, 2/3 and 4/5) were PCR amplified in our source material. The reasoning for selecting

these seven out of 14 introns in *nad* genes for further PCR analyses, which was based on outcomes of *in silico* PCR analyses and aimed at resolving obstacles evident from those analyses that may hamper PCR amplification and further utilization of these loci, is given in Results section upon providing outcomes of *in silico* PCR analysis for each out these seven loci.

We used mtDNA genome sequence of *V. faba* (NEGRUK, 2013) and FastPCR 6.3 trial version (KALENDAR *et al.*, 2009) to design new primers for PCR amplification of *nad1* intron 2/3 in this species. The difference in melting temperatures of F and R primers was set to 4 °C, CG clump was not opted, and primers of c. 20 bp length and CG content around 50% were generated. Out of several F and R primers which can be used for amplification of this particular intron, only those located within exon 2 (F primers) and exon 3 (R primers) and displaying quality above 75% were further used. Thus, we designed two F primers [*nad1e2f15* (5'-TCTAGGAGCATTACGATCTGCAG - 3') and *nad1e2f108* (5'-ATCCGCGAAGGCAATCGCTCG - 3')] and three R primers [*nad1e3r1484* (5'-AGTTTCCCTGCATGTGGCTCG - 3'), *nad1e3r1602* (5'-GGAACAAGGGAATACCGGACC - 3') and *nad1e3r1667* (5'-TGGGAGATCAAACGGAGCTCG - 3')]. Each out of two F primers was employed with each out of three R primers for PCR amplification of *nad1* intron 2/3 in *V. faba*.

PCR reactions were carried out in 25 µl volumes containing: 100 ng template DNA; 1 × *Taq* Buffer with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (Fermentas, Vilnius, Lithuania); 2.5 mM MgCl<sub>2</sub>; 0.2 mM dNTPs; 0.4 µM of each Forward (F) and Reverse (R) primer; 0.80 % BSA (Bovine Serum Albumin, Promega, St Louis, U.S.A.); and 0.025 U of *Taq* DNA polymerase (Fermentas, Vilnius, Lithuania). Reactions were performed in the Mastercycler Gradient machine (Eppendorf AG, Hamburg, Germany) under following conditions: denaturation at 95 °C for 5 min., 39 cycles of denaturation at 95 °C for 45 sec., annealing at temperatures 55 °C (*nad5* intron *d/e*, *nad7* introns 1/2 and 2/3), 58 °C (*nad1* intron 1/2, *nad4* introns 1/2 and 2/3, *nad7* intron 4/5) or 67 °C (used for all six combinations of new F and R primers for amplification of *nad1* intron 2/3), extension at 72 °C for 2 min., final extension at 72 °C for 10 min.. PCR products were subjected to electrophoresis through 2% agarose gels followed by ethidium-bromide staining and were visualised under the UV light utilizing BioDoc Analyze (Biometra, Göttingen, Germany).

PCR products obtained with new primers *nad1e2f15* and *nad1e3r1602* for amplification of *nad1* intron 2/3 were sequenced commercially with the F primer by Macrogen Europe, Amsterdam, Netherlands (<http://dna.macrogen.com/eng/>) via Sanger sequencing using 96-capillary 3730xl DNA Analyzer automated sequencer (Applied Biosystems, Inc., U.S.A.). Obtained sequences of the 5'-end of this intron in 16 *V. faba* genotypes were edited and aligned manually along with *V. faba* cultivar Broad Windsor sequence of this intron (NEGRUK, 2013) using Muscle (EDGAR, 2004) in MEGA 5.04 (TAMURA *et al.*, 2011).

## RESULTS

### *In silico* PCR analyses

The positions of all primer binding sites of 15 published primer pairs used for *in silico* PCR amplification of 14 cis-spliced introns in *V. faba*, *V. angularis* and *V. radiata*, the complementarity of primers and template DNA, *T<sub>m</sub>* of individuals primers, *T<sub>a</sub>* of primer pairs and lengths of obtained PCR products are presented in Table 1. With regard to the complementarity of primers and template DNAs of three legumes, *in silico* PCR analysis revealed that F and/or R primers from six primer pairs (for amplification of *nad2* intron 4/5, *nad4* introns 1/2, 3/4 and

primer pair of DEMESURE *et al.* (1995) for amplification of the entire *nad4* gene, *nad7* introns 2/3 and 3/4 harboured single nucleotide mismatches in their binding sites. Lower complementarity of primers and template DNAs was observed also for degenerate primers comprising up to three ambiguous nucleotides (F primers from primer pairs for amplification of *nad2* introns 1/2 and 3/4, and *nad7* intron 4/5).

Table 1 Results of *in silico* PCR amplification of 14 *cis*-spliced mitochondrial introns in *nad* genes in *Vicia faba*, *Vigna angularis* and *Vigna radiata*.

Species	F primer	5'-3'	Position	Com (%)	Tm (°C)	R primer	5'-3'	Position	Com (%)	Tm (°C)	Fr. len.	Ta (°C)
<b>Locus 1: <i>nad1</i> intron 2/3 (reference: 1, 3)</b>												
<i>V. faba</i>	<i>nad1</i> exon2	GCATTACGATCTGCAGCTCA	380141 ←	100	55	<i>nad1</i> exon3	GGAGCTCGATTAGTTTCTGC	378525 →	100	52,9	1636	60
			380160				378544					
	44926 ←	448329	100	52,9	1636	60						
	449945	178244 ←	65	9,2								
<i>V. angu-</i> <i>laris</i>	<i>nad1</i> exon2	GCATTACGATCTGCAGCTCA	222894 →	100	55	<i>nad1</i> exon3	GGAGCTCGATTAGTTTCTGC	224527 ←	100	52,9	1653	60
			222913				224546					
	35027 ←	493293 ←	65	9,2								
	35046	493312	65	9,2								
<i>V. radi-</i> <i>ata</i>	<i>nad1</i> exon2	GCATTACGATCTGCAGCTCA	3065 →	100	55	<i>nad1</i> exon3	GGAGCTCGATTAGTTTCTGC	4698 ←	100	52,9	1653	60
			3084				4717					
	179257 ←	202612 →	65	40,6								
	179276	202631	75	34,1								
<b>Locus 2: <i>nad1</i> intron 4/5 (reference: 2, 3) (note: contains <i>mark</i> gene)</b>												
<i>V. faba</i>	<i>nad1/4</i>	GCCAATATGATCTTAATGAG	89487 →	100	46,8	<i>nad1/5</i>	TCACCTTGATACTAAACCAG	92731 ←	100	48,4	3264	54
			89506				92750					
	4953 →	4972	70	22,1								
	457758 →	457777	70	22,1								
<i>V. angu-</i> <i>laris</i>	<i>nad1/4</i>	GCCAATATGATCTTAATGAG	67987 ←	100	46,8	<i>nad1/5</i>	TCACCTTGATACTAAACCAG	64738 →	100	48,4	3269	54
			68006				64757					
	65531 ←	62282 →	100	48,8	3269	54						
	65550	62301	100	48,8	3269	54						
<b>Locus 3: <i>nad2</i> intron 1/2 (reference: 3)</b>												
<i>V. faba</i>	<i>nad2/1</i>	AATGTGGGTGGCTTGGWTT	27785 →	98	54,6	<i>nad2/2</i>	AATATGTAAAATTGTCCTC	29101 ←	100	47,1	1336	53
			27804				29120					
	2331 ←	2350	78	34,7								
	99499 →	99518	62	41,6								
<i>V. angu-</i> <i>laris</i>	<i>nad2/1</i>	AATGTGGGTGGCTTGGWTT	455136 ←	98	54,6	<i>nad2/2</i>	AATATGTAAAATTGTCCTC	148978 ←	100	47,1	1374	53
			455155				148997					
	147624 →	147643	98	54,6								
	142527 →	142546	78	37,9								
<i>V. radi-</i> <i>ata</i>	<i>nad2/1</i>	AATGTGGGTGGCTTGGWTT	402176 →	68	17,5	<i>nad2/2</i>	AATATGTAAAATTGTCCTC	192568 →	100	47,1	1374	53
			402195				192587					
	193922 ←	193941	98	54,6								
	163456 ←	163475	78	37,9								
<b>Locus 4: <i>nad2</i> intron 3/4 (reference: 3)</b>												
<i>V. faba</i>	<i>nad2/3</i>	AGAAARGAATGCTGTAACCG	415414 →	98	50	<i>nad2/4</i>	ATGGGGATTIKTYARTATCGC		58	4	No product	
			415433									
	332368 ←	332387	70	40,7								
	152757 ←	152776	62	23,8								
<i>V. angu-</i> <i>laris</i>	<i>nad2/3</i>	AGAAARGAATGCTGTAACCG	99140 ←	98	50	<i>nad2/4</i>	ATGGGGATTIKTYARTATCGC		62	23,8	No product	
			99159									
	324667 ←	324686	62	23,8								
	267633 ←	267652	62	23,8								
<i>V. radi-</i> <i>ata</i>	<i>nad2/3</i>	AGAAARGAATGCTGTAACCG	96655 ←	98	50	<i>nad2/4</i>	ATGGGGATTIKTYARTATCGC		62	23,8	No product	
			96674									
	267633 ←	267652	62	23,8								
	267633 ←	267652	62	23,8								

Locus 5: <i>nad2</i> intron 4/5 (reference: 3)										
<i>V. faba</i>	<i>nad2/4</i>	TTCATATAGAAATCCATGTCC	413337 → 413356 139973 ← 139992	95 70	41,2 22,2	<i>nad2/5</i>	CTATTTGTCTTCGCCGCTT	414960 ← 414979	100	56,3 1643 54
<i>V. angu- laris</i>	<i>nad2/4</i>	TTCATATAGAAATCCATGTCC	101195 ← 101214	95	41,2	<i>nad2/5</i>	CTATTTGTCTTCGCCGCTT	356348 → 356367 99594 → 99613 89580 → 89599 97109 → 97128 87090 → 87109	70 100 75 100	28,6 56,3 29,3 56,3 1622 54
<i>V. radi- ata</i>	<i>nad2/4</i>	TTCATATAGAAATCCATGTCC	98711 ← 98730	95	41,2	<i>nad2/5</i>	CTATTTGTCTTCGCCGCTT	141750 → 141769 86387 ← 86406	70 100	33,1 53,9 2044 61
Locus 6: <i>nad4</i> intron 1/2 (reference: 1, 3)										
<i>V. faba</i>	<i>nad4 exon1</i>	CAGTGGGTGGTCTGGTATG	353611 → 353630	95	53,1	<i>nad4 exon2</i>	TCATATGGGCTACTGAGGAG	355634 ← 355653 418785 → 418804 88877 ← 88896	100 70	53,9 33,1 2043 61
<i>V. angu- laris</i>	<i>nad4 exon1</i>	CAGTGGGTGGTCTGGTATG	86853 → 86872 13644 → 13663	95 80	53,1 41,8	<i>nad4 exon2</i>	TCATATGGGCTACTGAGGAG	164233 ← 164252	70	33,1
<i>V. radi- ata</i>	<i>nad4 exon1</i>	CAGTGGGTGGTCTGGTATG	84363 → 84382 211303 → 211322	95 80	53,1 41,8	<i>nad4 exon2</i>	TCATATGGGCTACTGAGGAG		100	53,9 2044 61

Locus 7: <i>nad4</i> intron 2/3 (reference: 2, 3)										
<i>V. faba</i>	<i>nad4/2</i>	CTCCTCAGTAGCCCATATGA	355634 – >355653 82203 ← 82222	100 70	52,5 21,7	<i>nad4/3</i>	AACCAGTCCATGACTTAACA	358879 ← 358898	100	53,4 3265 60
<i>V. angu- laris</i>	<i>nad4/2</i>	CTCCTCAGTAGCCCATATGA	88877 → 88896 206274 → 206293 134816 ← 134835	100 75 70	52,5 22,7 21,7	<i>nad4/3</i>	AACCAGTCCATGACTTAACA	92197 ← 92216	100	53,4 3340 60
<i>V. radi- ata</i>	<i>nad4/2</i>	CTCCTCAGTAGCCCATATGA	86387 → 86406 132278 ← 132297	100 70	52,5 21,7	<i>nad4/3</i>	AACCAGTCCATGACTTAACA	89708 ← 89727	100	53,4 3341 60
Locus 8: <i>nad4</i> intron 3/4 (reference: 2, 3)										
<i>V. faba</i>	<i>nad4 exon3</i>	GGAGCTTCCAAGAAATAG	359082 – > 359101 319535 – > 319554	100 65	49,7 16,3	<i>nad4 exon4</i>	GCCATGTTGCACTAAGTTAC	362092 ← 362111	100	52,8 3030 57
<i>V. angu- laris</i>	<i>nad4 exon3</i>	GGAGCTTCCAAGAAATAG	92400 → 92419 125508 – > 125527	100 75	49,7 34,9	<i>nad4 exon4</i>	GCCATGTTGCACTAAGTTAC	38312 → 38331 95180 ← 95199	70 95	34,5 50,6 2800 57
<i>V. radi- ata</i>	<i>nad4 exon3</i>	GGAGCTTCCAAGAAATAG	89911 → 89930	100	49,7	<i>nad4 exon4</i>	GCCATGTTGCACTAAGTTAC	92694 ← 92713 122970 → 122989	95 75	50,6 2803 35,6 57
Locus 9: <i>nad4</i> subunit (reference: 1) (note: F primer is located within exon1 of <i>nad4</i> gene)										
<i>V. faba</i>	<i>nad4 exon2</i>	TGTTCCCGAAGCGACACTT	355516 → 355535	95	52	<i>nad4 exon4</i>	GGAACACTTTGGGGTGAACA	362048 ← 362067	100	53,2 6552 64
<i>V. angu- laris</i>	<i>nad4 exon2</i>	TGTTCCCGAAGCGACACTT	88759 → 88778	95	52	<i>nad4 exon4</i>	GGAACACTTTGGGGTGAACA	95136 ← 95155	100	53,2 6397 64
<i>V. radi- ata</i>	<i>nad4 exon2</i>	TGTTCCCGAAGCGACACTT	86269 → 86288	95	52	<i>nad4 exon4</i>	GGAACACTTTGGGGTGAACA	92650 ← 92669	100	53,2 6401 64

Locus 10: <i>nad5</i> intron a/b (reference: 2, 3)										
<i>V. faba</i>	<i>nad5/1</i>	TTTTTCGGGACGTTTCTAG	71584 →			<i>nad5/2</i>	TTTGGCCAAGTATCCTACAA	73834 ←		
			71603	100	49,1			73853	100	50,3 2270 57
<i>V. angu-</i>	<i>nad5/1</i>	TTTTTCGGGACGTTTCTAG	120992 →			<i>nad5/2</i>	TTTGGCCAAGTATCCTACAA	118744 ←		
<i>laris</i>			121011	100	49,1			118763	100	50,3 2268 57
<i>V. radi-</i>	<i>nad5/1</i>	TTTTTCGGGACGTTTCTAG	118473 →			<i>nad5/2</i>	TTTGGCCAAGTATCCTACAA	116224 ←		
<i>ata</i>			118492	100	49,1			116243	100	50,3 2269 57
Locus 11: <i>nad5</i> intron d/e (reference: 2, 3)										
<i>V. faba</i>	<i>nad5/4</i>	CCAATTTTGGGCAATTC	279736 →			<i>nad5/5</i>	CATTGCAAAGGCATAATGAT	281047 ←		
			279755	100	55,4			281066	100	49,2 1331 57
			563870 →					562559 →		
			563889	100	55,4			562578	100	49,2 1331 57
			309974 ←							
			309993	70	29,4					
			367879 ←							
			367898	70	29,4					
			437664 ←							
			437683	70	29,4					
<i>V. angu-</i>	<i>nad5/4</i>	CCAATTTTGGGCAATTC	183939 →			<i>nad5/5</i>	CATTGCAAAGGCATAATGAT	185255 ←		
<i>laris</i>			183958	100	55,4			185274	100	49,2 1336 57
								367370 →		
								367389	70	26,4
								401251 →		
								401270	70	20,2
<i>V. radi-</i>	<i>nad5/4</i>	CCAATTTTGGGCAATTC	324793 ←			<i>nad5/5</i>	CATTGCAAAGGCATAATGAT	323477 →		
<i>ata</i>			324812	100	55,4			323496	100	49,2 1336 57
								35055 →		
								35074	70	26,1
								292633 →		
								292652	70	20,2

Locus 12: <i>nad7</i> intron 1/2 (reference: 2, 3) (note: R primers generate non-specific PCR product of 3244 bp in <i>V. angularis</i> , <i>Ta</i> = 56°C)										
<i>V. faba</i>	<i>nad7/1</i>	ACCTCAACATCCTGCTGCTC	226324 ←			<i>nad7/2</i>	CGATCAGAATAAGGTAAAGC	225275 →	100	48,8 1069 55
			226343	100	59,2			225294		
			302233 →					303282 ←		
			302252	100	59,2			303301	100	48,8 1069 55
			541373 ←					540324 →		
			541392	100	59,2			540343	100	48,8 1069 55
								34233 →		
								34252	75	11,3
								94184 →		
								94203	70	27,9
<i>V. angu-</i>	<i>nad7/1</i>	ACCTCAACATCCTGCTGCTC	356308 →			<i>nad7/2</i>	CGATCAGAATAAGGTAAAGC	357341 ←		
<i>laris</i>			356327	100	59,2			357360	100	48,8 1053 55
								354117 →		
								354136	75	27,9
								199933 ←		
								199952	70	13,5
								115191 →		
								115210	65	22,8
								266002 ←		
								266021	75	11,3
<i>V. radi-</i>	<i>nad7/1</i>	ACCTCAACATCCTGCTGCTC	343682 →			<i>nad7/2</i>	CGATCAGAATAAGGTAAAGC	344715 ←		
<i>ata</i>			343701	100	59,2			344734	100	48,8 1035 49
								308812 →		
								308831	70	13,5
								284540 →		
								284559	75	24,9
								112666 →		
								112685	65	22,8
								141365 →		
								141384	75	11,3
								126095 ←		
								126114	65	16,5

Locus 14: <i>nad7</i> intron 3/4 (reference: 2, 3)										
<i>V. faba</i>	<i>nad7/3</i>	TCTATGATGGCCCAAGAACA	223919 ←		<i>nad7/4</i>	ACACCAAATTCCTTTAGG	220057 →			
			223938	95	48,7		220076	100	52,5	3882 58
			304638 →				308500 ←			
			304657	95	48,7		308519	100	52,5	3882 58
			538968 ←				535106 →			
			538987	95	48,7		535125	100	52,5	3882 58
							436190 ←			
							436209	100	52,5	
							366405 ←			
							366424	100	52,5	
<i>V. angu- laris</i>	<i>nad7/3</i>	TCTATGATGGCCCAAGAACA	358711 →		<i>nad7/4</i>	ACACCAAATTCCTTTAGG	362576 ←			
			358730	95	48,7		362595	100	52,5	3885 58
			266477 ←							
			266496	65	35,9					
							214018 ←			
							214037	80	16,4	
							349940 ←			
							349959	100	52,5	3874 58
<i>V. radi- ata</i>	<i>nad7/3</i>	TCTATGATGGCCCAAGAACA	346086 →		<i>nad7/4</i>	ACACCAAATTCCTTTAGG				
			346105	95	48,7					
			140890 →							
			140909	65	35,9					
							395462 ←			
							395481	75	7,4	

Locus 13: <i>nad7</i> intron 2/3 (reference: 2, 3) (note: F primers generate non-specific PCR product of 3990 bp in <i>V. faba</i> , <i>Ta</i> = 56°C)										
<i>V. faba</i>	<i>nad7/2</i>	GCTTACCTTATTCTGATCG	225275 ←		<i>nad7/3</i>	TGTTCTTGGCCATCATAGA	223919 →			
			225294	100	49,3		223938	95	49,8	1376 55
			303282 →				304638 ←			
			303301	100	49,3		304657	95	49,8	1376 55
			540324 ←				538968 →			
			540343	100	49,3		538987	95	49,8	1376 55
			221305 →							
			221324	70	17,9					
			307252 ←							
			307271	70	17,9					
			536354 →							
			536373	70	17,9					
<i>V. angu- laris</i>	<i>nad7/2</i>	GCTTACCTTATTCTGATCG	357341 ←		<i>nad7/3</i>	TGTTCTTGGCCATCATAGA	358711 ←			
			>357360	100	49,3		358730	95	49,8	1390 55
			344425 →							
			344444	75	12,3					
			324826 →							
			324845	70	23,3					
<i>V. radi- ata</i>	<i>nad7/2</i>	GCTTACCTTATTCTGATCG	344715 →		<i>nad7/3</i>	TGTTCTTGGCCATCATAGA	346086 ←			
			344734	100	49,3		346105	95	49,8	1391 55
			231076 →							
			231095	75	12,3					
			267792 →							
			267811	70	23,3					

Locus 15: <i>nad7</i> intron 4/5 (reference: 3)										
<i>V. faba</i>	<i>nad7/4</i>	TGTCCTCCATCACGATVTCG	222223 ←		<i>nad7/5</i>	CCAAATTCCTTTAGGTGC	220060 →			
			222242	97	52,8		220079	100	51,8	2183 60
			306334 →				308497 ←			
			306353	97	52,8		308516	100	51,8	2183 60
			537272 ←				535109 →			
			537291	97	52,8		535128	100	51,8	2183 60
			17125 →							
			17144	72	13,3					
			67213 →							
			6723272	72	13,3					
							366402 ←			
							366421	100	51,8	
							436187 ←			
							436206	100	51,8	
<i>V. angu- laris</i>	<i>nad7/4</i>	TGTCCTCCATCACGATVTCG	360433 →		<i>nad7/5</i>	CCAAATTCCTTTAGGTGC	362573 ←			
			360452	97	52,8		362592	100	51,8	2160 60
							59241 ←			
							59260	70	25,3	
<i>V. radi- ata</i>	<i>nad7/4</i>	TGTCCTCCATCACGATVTCG	347810 →		<i>nad7/5</i>	CCAAATTCCTTTAGGTGC	349937 ←			
			347829	97	52,8		349956	100	51,8	2147 60
							56773 ←			
							56792	70	25,3	

References: 1 – DEMESURE *et al.* (1995); 2 – DUMOLIN-LAPEGUE *et al.* (1997); 3 – DUMINIL *et al.* (2002); Com – complementarity of a primer and template DNA expressed through the percentage of binding nucleotides; *T<sub>m</sub>* – melting temperature of a primer in °C; Fr. len. – the length of the PCR product obtained via *in silico* PCR method; *T<sub>a</sub>* – annealing temperature of a primer pair in °C.

Primer pair for amplification of the *nad2* intron 3/4 failed to generate PCR product in all three legumes because its R primer, which comprised three ambiguous nucleotides, lacked primer binding sites within exon4 of the *nad2* gene, and primed loosely (complementarity between 58% and 62%) to alternative locations in all taxa. The remaining 14 primer pairs generated desired PCR products in all three legumes and their sizes ranged from c. 1 kb (*nad7* intron 1/2) to 3.9 kb (*nad7* intron 3/4), while the length of the PCR product obtained with primers F: *nad4* exon2 and R: *nad4* exon4 of DEMESURE *et al.* (1995) was c. 6.5 kb in all three legumes. We found that F primer from this primer pair was located not within exon2 but within exon1.

Out of 13 primer pairs that generated desired PCR products long up to 4 kb, only F and R primers from a primer pair for amplification of the *nad5* intron a/b displayed single binding sites with maximal complementarity with template DNAs of all three species. Many primers from the remaining 12 primer pairs, however, had not only highly complementary binding sites at expected locations within mtDNA genomes of studied legumes, but also additional (multiple) binding sites of lower complementarity (between 65% and 98%). We tested via PCR amplification in 16 genotypes of *V. faba* whether multiple binding sites of primers would lead to the occurrence of multiple amplification products.

The highest number of multiple binding sites was observed for R primer from a primer pair for amplification of the *nad7* intron 1/2 in all legumes (up to five additional binding sites of lower complementarity). On the other hand, the entire *nad7* gene was triplicated within *V. faba* mtDNA genome and exon5 of this gene was copied on two alternative locations (NEGRUK, 2013). Also, loci *nad1* intron 2/3 and *nad5* intron d/e were duplicated in this species (NEGRUK, 2013) and R and F primer from a primer pair for amplification of the former and the latter locus, respectively, also displayed additional binding sites of lower complementarity. We PCR amplified above listed loci (*nad1* intron 2/3, *nad5* intron d/e, *nad7* introns 1/2 and 4/5) to test not only the potential confounding effect of the presence of two or three copies of introns on their utilization for further manipulation but also the effect of additional binding sites with maximal or lower complementarity with template DNA on occurrence of multiple amplification products.

Along with desired PCR products, non-specific amplification products were obtained via *in silico* PCR at two loci, *nad7* intron 2/3 in *V. faba* (c. 4 kb) and *nad7* intron 1/2 in *Vigna angularis* (c. 3.3 kb). They were generated with F primers (at the former locus) and with R primers (at latter locus). The occurrence of these multiple amplification products predicted via *in silico* PCR was tested via PCR amplification in 16 genotypes of *V. faba*.

> Table 1

#### *PCR amplification of seven introns in nad genes in V. faba*

Based on nucleotide content of mtDNA sequences of studied legumes comprising binding sites of primers, we replaced mispairing nucleotides in three primers (F primer from a primer pair for amplification of the *nad7* intron 4/5 – we introduced nucleotide G at position 17 instead of an ambiguous nucleotide V; F primer from a primer pair for amplification of the *nad4* intron 1/2 - T at first position instead of C; and R primer from a primer pair for amplification of *nad7* intron 2/3 -

G at position 16 instead of A), and used such improved primers for PCR amplification in 16 genotypes of *V. faba*.

The outcomes of PCR amplification of seven introns in *nad* genes in 16 genotypes of *V. faba* are presented in Figure 1 (a to g). Surprisingly, PCR amplification of *nad1* intron 2/3 was rather poor because PCR products of expected length (c. 1.6 kb) were obtained in five out of 16 individuals, while in other individuals, either faint bands of expected length or lack of PCR products were detected (Figure 1a). Furthermore, smears, which may occur due to various reasons including high DNA concentrations, low annealing temperatures, mispriming caused by secondary structure template, etc., were persistent even with altered DNA concentrations and PCR conditions (data not shown). Faint bands of expected length were obtained in one cultivar at locus *nad5* intron d/e (Rojo, Peru) and in two cultivars at locus *nad7* intron 4/5 (BPL4225, Morocco, and Diana, Germany), while lack of PCR products was observed in one cultivar at each out of two loci (Deligrad, Serbia at locus *nad4* intron 2/3, and Rojo, Peru at locus *nad7* intron 1/2). Overall, single bands of expected length suitable for further manipulation were obtained at three loci, *nad4* intron 2/3, *nad7* introns 2/3 and 4/5 (Figure 1c, f and g, respectively), while multiple amplification products were obtained at three loci, *nad4* intron 1/2, *nad5* intron d/e and *nad7* intron 1/2 (Figure 1b, d and e, respectively). At locus *nad7* intron 4/5, variation in length in 16 genotypes of *V. faba* was evident even on agarose gels (Figure 1g).

PCR amplification of *nad1* intron 2/3 with PCR primers provided in this study (two F and three R primers) was successful, and only outcomes of PCR amplification with *nad1e2f15* and *nad1e3r1602* are presented in Figure 1h.

Readable electropherograms of PCR products in 16 *V. faba* genotypes obtained with primers *nad1e2f15* and *nad1e3r1602* and sequenced with F primer were of c. 700 bp, and were alignable among themselves and with the 5'-end of this intron from *V. faba* cultivar Broad Windsor whose full length is 1439 bp (NEGRUK, 2013). Within the 5'-end of the intron, two transversions were observed while length mutations were lacking. With regard to the primary nucleotide sequence of this intron reported by NEGRUK (2013), T/G transversion was detected at position 128, and A/C transversion at position 236.

## DISCUSSION

A burst of complete mtDNA sequences of flowering plants especially over the past few years is essential for facilitating broader utilization of this haploid and predominantly uniparentally-inherited genome in diverse studies in plants. This is because available mtDNA sequences of taxa of interest or those of their close relatives may be used for detection of regions displaying variability in primary nucleotide sequences or in length, as well as for delineation of more conserved regions suitable for designing primers for PCR amplification of those informative regions. However, although mtDNA genomes of higher plants have rather slow evolution of primary nucleotide sequences (WOLFE *et al.*, 1987; LAROCHE *et al.*, 1997), they are exceptionally variable in size and organization and are prone to frequent insertions, deletion and structural rearrangements (LAROCHE *et al.*, 1997; NEGRUK, 2013 and references therein). Due to such fluid nature of plant mtDNA genomes, it is possible that mtDNA sequences available in public databases may differ to the certain extent from those of studied individuals and thus, the predictions based on known mtDNA sequences may not be fully applicable to unknown sequences.




Figure 1 Results of PCR amplification of seven introns in *nad* genes in 16 genotypes of *Vicia faba* representative of the worldwide gene pool of this species; a to g – PCR products obtained with published original or improved PCR primers; h - PCR products obtained with primers developed in this study (*nad1e2f15* and *nad1e3r1602*) for amplification of *nad1* intron 2/3.

A negative image of an ethidium bromide-stained agarose gel is shown. The size markers L1 and L2 in the lanes 1 and 10, respectively, are fragments of a 1 kb DNA ladder and 50 DNA bp ladder; reference bands of the size marker L2 correspond to fragments of 0.25 kb, 0.5 kb and 1 kb, respectively; PCR products in *Vicia faba* traditional cultivars (TC) or improved cultivars (IC) of *major* (Ma), *minor* (Mi) and *equina* (Eq) type and subsp. *paucijuga* (Pau) are presented in lanes 2 - 8 and 11 - 18, and comprise cultivars: 1 - Troy (country of origin: Denmark, characteristics: Eq, IC); 2 - CH170 (United Kingdom - Eq, IL); 3 - E14b (Ethiopia - Eq, TC); 4 - Mikko (Finland - Mi, TC); 5 - Apsalos (Greece - Eq, IL); 6 - CH182 (India - Pau, TC); 7 - Violetta di policoro (Italy - Ma, TC); 8 - Kingpi (China - Eq, TC); 9 - BPL4225 (Morocco - not available); 10 - Diana (Germany - Mi, IC); 11 - Silian (Sudan - Mi, TC); 12 - Feve Seville (Spain - Ma, TC); 13 - Rojo Peru Ma, TC); 14 - Bački Petrovac 2 (Serbia - Ma, TC); 15 - Deligrad (Serbia - Ma, TC); and 16 - 159B (former Yugoslavia - Mi, TC).

In this study, we utilized both predictions based on *in silico* PCR analyses and PCR amplification in our source material in order to provide a set of PCR primers suitable for amplification of potentially informative introns in *nad* genes in *V. faba* which would facilitate broader utilization of mtDNA genome in phylogeographic and other studies in this species for which unambiguous data on wild ancestors, origin and domestication are lacking.

#### *In silico* PCR analyses

*In silico* PCR analyses revealed that six out of 15 tested universal primer pairs of DEMESURE *et al.* (1995), DUMOLIN-LAPEGUE *et al.* (1997) and DUMINIL *et al.* (2002) used for predictions of amplification of all 14 *cis*-spliced introns in *nad* genes in mitochondrial genomes of *V. faba* and two *Vigna* species were not suitable for these species because they harboured mispaired nucleotides within their binding sites. Although this obstacle can be overcome, this questions universality of those primers which, however, may be increased via introduction of ambiguous nucleotides. On the other hand, we found that utilization of degenerate primers harbouring ambiguous nucleotides in primers from three out of 15 primer pairs was not necessary in studied legumes. Furthermore, we argue that in legumes, characterized by large and repetitive mtDNA genomes (e.g. the size of mtDNA genome in *V. faba* was estimated to 588 kb, NEGRUK, 2013; 404466 bp in *Vigna angularis*, NAITO *et al.*, 2013; and 401262 bp in *Vigna radiata*, ALVERSON *et al.*, 2011), utilization of degenerate primers should be avoided because they may either fail to generate PCR product (e.g. at locus *nad2* intron 3/4), or alternatively, may display additional (multiple) binding sites with lower complementarity with template DNAs that may lead to the occurrence of multiple amplification products in PCR reactions (e.g. at loci *nad4* intron 1/2, *nad5* intron d/e and *nad7* intron 1/2).

With the exception of a primer pair for amplification of the *nad2* intron 3/4, the remaining primer pairs generated desired PCR products of up to c. 4 kb in three legume species, while the length of the PCR product obtained with primers F: *nad4* exon2 and R: *nad4* exon4 of DEMESURE *et al.* (1995) was c. 6.5 kb in all three legumes. DEMESURE *et al.* (1995) reported that these primers generated a PCR product of c. 4 kb in eight plant species. However, we found that such a long products were obtained because F primer was located not within exon2 but within exon1, and when employed with the R primer located within exon4, amplified the entire *nad4* gene comprised of four exons and three introns. Such a long PCR products, although suitable for studies with restriction enzymes (DEMESURE *et al.*, 1995), are rather challenging for the assessment of their primary nucleotide sequences via sequencing reactions. For instance, even mtDNA stretches of c. 2 kb have been PCR amplified via multiple PCR reactions whose products have been used to assess the nucleotide content of the entire 2 kb long region (e.g. HAVANANDA *et al.*, 2010). In *V. faba*, multiple PCR reactions or sequencing with internal primers may be required for seven loci whose lengths estimated via *in silico* PCR analyses were 2 kb to 3.9 kb (*nad1* intron 4/5, *nad4* introns 1/2, 2/3 and 3/4, *nad5* intron a/b and *nad7* introns 3/4 and 4/5).

Overall, out of 15 published universal primer pairs of DEMESURE *et al.* (1995), DUMOLIN-LAPEGUE *et al.* (1997) and DUMINIL *et al.* (2002), *in silico* PCR analyses revealed that only one (for amplification of the *nad5* intron a/b) is suitable for utilization in *V. faba* and representatives of the genus *Vigna*. Primers from this primer pair displayed single maximally complementary binding sites in all studied species and generated PCR products of c. 2.3 kb suitable for further manipulation. The main obstacles hampering utilization of other primer pairs comprised generation of exceptionally long PCR products and multiple binding sites which may lead to the

occurrence of multiple amplification products. However, as already mentioned, predictions obtained via *in silico* PCR method may not necessarily be concordant with patterns obtained via PCR reactions, as discussed below.

#### *PCR amplification of seven introns in nad genes in V. faba*

The most unexpected outcome of PCR amplification of seven introns in *nad* genes in 16 genotypes of *V. faba* was poor amplification of locus *nad1* intron 2/3 with universal primers of DEMESURE *et al.* (1995). This is because this locus was successfully PCR amplified with same primers in 13 representatives of the genus *Vicia* including *V. faba* (RYZHOVA *et al.*, 2012). Nonetheless, *nad1* intron 2/3 was successfully PCR amplified in our material as well but with primers reported in this study.

PCR amplification of other six loci in our material revealed that intron 2/3 of the *nad4* gene and introns 2/3 and 4/5 of the *nad7* gene (the latter two being triplicated within mtDNA genome of *V. faba*, NEGRUK, 2013) are also suitable for future studies in this species because at these loci, multiple amplification products were lacking and PCR products of expected length were generated in all individuals with one exception only (Deligrad, Serbia at locus *nad4* intron 2/3). Thus, it appears that duplications and triplications of mtDNA introns within mtDNA genome of *V. faba* may not represent an obstacle for their utilization. This is also supported by comparison of primary nucleotide sequences of all duplicated and triplicated mtDNA introns in *V. faba* cultivar Broad Windsor because all corresponding copies of introns harboured identical nucleotide content (data not shown). Although further work is required to reveal whether these three loci are informative for future studies in *V. faba*, variability in length has been observed at *nad7* intron 4/5 even on agarose gels (Figure 1g). We assume that length variation at this locus was not related to the presence of five maximally complementary binding sites of the R primer from a primer pair for amplification of this locus.

Loci *nad4* intron 1/2, *nad5* intron d/e and *nad7* intron 1/2 are not recommended for future utilization in *V. faba* because primers that amplify these loci are likely to generate not only desired PCR products of expected length but also additional non-specific amplification products of various lengths. Such an outcome, however, was not fully concordant with predictions on occurrence of multiple amplification products obtained in *in silico* analyses which revealed multiple amplification products at locus *nad7* intron 2/3 only. Nonetheless, multiple binding sites of primers within mtDNA genome of *V. faba* cultivar Broad Windsor, which signify the presence of several oligonucleotide stretches within mtDNA genome of this individual that are more or less complementary with the primary nucleotide sequence of utilized primers and may contribute towards the occurrence of multiple amplification products, were observed for majority of primers used in this study. For instance, at locus *nad4* intron 1/2, *in silico* PCR revealed only one additional binding site of R primer used for amplification of this locus, while in PCR reaction, the highest number of non-specific bands was observed at this locus (Figure 1b).

The observed discordance with regard to the occurrence of multiple amplification products which were predicted via *in silico* analyses and obtained in PCR reactions may be due to the more technical issues, such as limitations of the utilized *in silico* method which may or may not be able to delineate all more or less complementary binding sites of primers that may generate multiple amplification products, or alternatively, to the differences in organization and/or primary nucleotide sequences of mtDNA genomes used for *in silico* analyses and those used for PCR amplification. The latter is possible owing to the exceptionally fluid organization of mtDNA

genomes in plants (LAROCHE *et al.*, 1997; NEGRUK, 2013 and references therein) and potential hampering effects of heteroplasmy (WOLOSZYNSKA, 2010). Thus, in case that our plant material used for PCR amplification was also used for *in silico* PCR analyses, it is possible that additional binding sites of primers, which generated multiple amplification products at loci *nad4* intron 1/2, *nad5* intron d/e and *nad7* intron 1/2, may have been detected. Therefore, we suggest that *in silico* PCR method, which utilizes mtDNA sequences that may differ to the certain extent from those of studied taxa, is not only suitable but also recommended for overcoming obstacles with regard to the mispairing of nucleotides within primer binding sites especially when based on mtDNA genome sequences of several related taxa. *In silico* PCR method is also suitable for estimating the length of PCR products, while its utility in predicting the occurrence of multiple amplification products is limited.

Finally, we support previous reports on intraspecific variability at the mtDNA level in *V. faba* (SCALLAN and HARMEY, 1996) because variability in length was detected at *nad7* intron 4/5, while variability in primary nucleotide sequence was found within the 5'-end of the second intron of the *nad1* gene based on analyses in 16 genotypes of *V. faba* representative of the worldwide gene pool of this species. The latter locus is one of the most commonly employed non-coding mtDNA regions in various studies in higher plants (e.g. FREUDENSTEIN and CHASE, 2001; WON and RENNER, 2003; ZHANG *et al.*, 2006; TOLLEFSRUD *et al.*, 2009; ALEKSIĆ and GEBUREK, 2010; HAVANANDA *et al.*, 2010; RYZHOVA *et al.*, 2012; LOCKWOOD *et al.*, 2013) and its future utilization in *V. faba* requires PCR amplification with primers provided in this study.

#### *Mitochondrial primers for amplification of introns in nad genes in V. faba*

The aim of our study was to provide molecular tools for broader utilization of variability of the mtDNA genome in phylogeographic, phylogenetic and other studies in *V. faba* that may shed more light on its origin, domestication and evolutionary relations. We focused primarily on introns in *nad* genes because non-coding mtDNA regions are evolving faster than their adjacent coding regions, and, as mentioned above, introns in *nad* genes are amongst the most commonly used mtDNA regions in various studies in higher plants.

Out of 15 universal primer pairs published by DEMESURE *et al.* (1995), DUMOLIN-LAPEGUE *et al.* (1997) and DUMINIL *et al.* (2002) which amplify all 14 *cis*-spliced introns in *nad* genes, only one (for amplification of *nad5* intron a/b) is suitable for *V. faba*, as inferred from *in silico* analyses. Based on PCR amplification of seven introns in 16 genotypes of *V. faba* originating from Europe, Africa, Asia and south America which represent all botanical varieties of this species (*major*, *minor*, *equina* and subsp. *paucijuga*) of various levels of improvement (traditional and improved cultivars), however, either original or improved primers from three primer pairs (for amplification of *nad4* intron 2/3 and *nad7* introns 2/3 and 4/5) generated desired PCR products suitable for further manipulation. Along with PCR primers for amplification of *nad1* intron 2/3 provided in this study, we report a set of primers suitable for PCR amplification of five introns in *nad* genes in *V. faba*. Although further studies are required to reveal whether these introns are informative, variability in primary nucleotide sequence was observed within 5'-end of the *nad1* intron 2/3, while length variability was detected at locus *nad7* intron 4/5. We also suggest further efforts to infer whether the remaining introns in *nad* genes that were not PCR amplified in this study may be utilized in *V. faba* as well, and also testing of other non-coding mtDNA regions for potential utilization in this species.

## ACKNOWLEDGEMENTS

Research grant 173005 of the Ministry of Education, Science and Technological Development of the Republic of Serbia, "Pavle Savić" program of bilateral collaboration between Republic of Serbia and Republic of France for 2012-2013, project number 680-00-132/2012-09/15.

Received April 20<sup>th</sup>, 2014

Accepted September 05<sup>th</sup>, 2014

## REFERENCES

- AKIBODE, S., M. MAREDA (2011): Global and regional trends in production, trade and consumption of food legume crops. Grad Res Masters Degree Plan B Papers, 114247. Department of Agricultural, Food, and Resource Economics, East Lansing, USA: Michigan State University Press, pp 83.
- ALEKSIĆ, J.M., T. GEBUREK (2010): Mitochondrial DNA reveals complex genetic structuring in a stenoendemic conifer *Picea omorika* [(Panč.) Purk.] caused by its long persistence within the refugial Balkan region. *Plant Syst Evol* 285(1-2): 1-11.
- ALEKSIĆ, J.M., D. STOJANOVIĆ, B. BANOVIĆ, R. JANČIĆ (2012): A simple and efficient DNA isolation method for *Salvia officinalis*. *Biochem Genet* 50(11-12): 881 – 892.
- ALVERSON, A.J., S. ZHUO, D.W. RICE, D.B. SLOAN, J.D. PALMER (2011): The mitochondrial genome of the legume *Vigna radiata* and the analysis of recombination across short mitochondrial repeats. *PLoS ONE* 6: e16404.doi:10.1371/journal.pone.0016404.
- AVISE, J.C., J. ARNOLD, R.M. BALL, E. BERMINGHAM, T. LAMB, J.E. NEIGEL, C.A. REEB, N.C. SAUNDERS (1987): Intraspecific phylogeography: the mitochondrial DNA bridge between population genetics and systematics. *Annu Rev Ecol Syst* 18: 489-522.
- AVISE, J.C. (2009): Phylogeography: retrospect and prospect. *J Biogeogr* 36(1): 3-15.
- CUBERO, J.I. (1974): On the evolution of *Vicia faba*. *Theor Appl Genet* 45(2): 47-51.
- DEMASURE, B., N. SODZI, R.J. PETIT (1995): A set of universal primers for amplification of polymorphic non-coding regions of mitochondrial and chloroplast DNA in plants. *Mol Ecol* 4(1): 129-134.
- DUC, G., S. BAO, M. BAUM, B. REDDEN, M. SADIKI, M.J. SUSO, M. VISHNIAKOVA, X. ZONG (2010): Diversity maintenance and use of *Vicia faba* L. genetic resources. *Field Crop Res* 115(3): 270-278.
- DUMINIL, J., M.H. PEMONGE, R.J. PETIT (2002): A set of 35 consensus primer pairs amplifying genes and introns of plant mitochondrial DNA. *Mol Ecol Notes* 2: 428-430.
- DUMOLIN-LAPEGUE, S., M.H. PEMONGE, R.J. PETIT (1997): An enlarged set of consensus primers for the study of organelle DNA in plants. *Mol Ecol* 6: 393-397.
- EDGAR, R.C. (2004): MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32(5): 1792-1797.
- FENNELL, S.R., W. POWELL, F. WRIGHT, G. RAMSAY, R. WAUGH (1998): Phylogenetic relationships between *Vicia faba* (Fabaceae) and related species inferred from chloroplast trnL sequences. *Plant Syst Evol* 212(3-4): 247-259.
- FREUDENSTEIN, J.V., M.W. CHASE (2001): Analysis of mitochondrial *nad1b-c* intron sequences in Orchidaceae: utility and coding of length-change characters. *Syst Bot* 26(3): 643-657.
- FROELICHER, Y., W. MOUHAYA, J.B. BASSENE, G. COSTANTINO, M. KAMIRI, F. LURO, R. MORILLON, P. OLLITRAULT (2011): New universal mitochondrial PCR markers reveal new information on maternal citrus phylogeny. *Tree Genet Genomes* 7(1): 49-61.
- HAIDER, N, I. NABULSI, N. MIRALI (2012): Identification of species of *Vicia* subgenus *Vicia* (Fabaceae) using chloroplast DNA data. *Turk J Agric For* 36: 297-308.

- HAVANANDA, T., E.C. BRUMMER, I.J. MAUREIRA-BUTLER, J.J. DOYLE (2010): Relationships among diploid members of the *Medicago sativa* (Fabaceae) species complex based on chloroplast and mitochondrial DNA sequences. *Syst Bot* 35(1): 140-150.
- KALENDAR, R., D. LEE, A.H. SCHULMAN (2009). FastPCR software for PCR primer and probe design and repeat search. *G3* 3: 1-14.
- LAROCHE, J., P. LI, L. MAGGIA, J. BOUSQUET (1997): Molecular evolution of angiosperm mitochondrial introns and exons. *P Natl Acad Sci USA* 94(11): 5722-5727.
- LOCKWOOD, J.D., J.M. ALEKSIĆ, J. ZOU, J. WANG, J. LIU, S.S. RENNER (2013): A new phylogeny for the genus *Picea* from plastid, mitochondrial, and nuclear sequences. *Mol Phylogenet Evol* 69(3): 717-727.
- MAXTED, N. (1993): A phenetic investigation of *Vicia* L. subgenus *Vicia* (Leguminosae Viciae). *Bot J Linn Soc* 111(2): 155-182.
- MAXTED, N. (1995): An ecogeographical study of *Vicia* subgenus *Vicia*. *Systematic and Ecogeographical Studies on Crop Genepools*, vol 8. IPGRI, Rome.
- MAXTED, N., M.A. CALLIMASSIA, M.D. BENNET (1991): Cytotaxonomic studies of Eastern Mediterranean *Vicia* species (Leguminosae). *Plant Syst Evol* 177(3-4): 221-234.
- MURATOVA, V.S. (1931): Common beans (*Vicia faba* L.). *B Appl Bot Genet Plant Breed Suppl* 50: 1-298.
- MOGENSEN, H.L. (1996): The hows and whys of cytoplasmic inheritance in seed plants. *Am J Bot* 83(3): 383-404.
- NAITO, K., A. KAGA, N. TOMOOKA, M. KAWASE (2013): De novo assembly of the complete organelle genome sequences of azuki bean (*Vigna angularis*) using next-generation sequencers. *Breed Sci* 63(2): 176-182.
- NEGRUK, V. (2013): Mitochondrial genome sequence of the legume *Vicia faba*. *Front Plant Sci* 4: 1-11.
- POTOKINA, E., N. TOMOOKA, D.A. VAUGHAN, T. ALEXANDROVA, R.Q. XU (1999): Phylogeny of *Vicia* subgenus *Vicia* (Fabaceae) based on analysis of RAPDs and RFLP of PCR-amplified chloroplast genes. *Genet Resour Crop Evol* 46: 149-161.
- RYZHOVA, N.N., E.A. D'ACHENKO, M.A. VISHNYAKOVA, E.Z. KOCHIEVA (2012): Characterization of variability of the gene *nad1* intron in representatives of *Vicia* L. genus (Fabaceae Lindl family). *Russ J Genet Appl Res* 2(2): 152-159.
- SCALLAN, C.D., M.A. HARMEY (1996): The use of mitochondrial DNA polymorphisms to distinguish commercial cultivars of the broad bean, *Vicia faba* L. *Euphytica* 87(3): 225-235.
- SCHAEFER, H., P. HECHENLEITNER, A. SANTOS-GUERRA, M.M. DE SEQUEIRA, R.T. PENNINGTON, G. KENICER, M.A. CARINE (2012): Systematics, biogeography, and character evolution of the legume tribe Fabeae with special focus on the middle-Atlantic island lineages. *BMC Evol Biol* 12: 250.
- SHIRAN, B., A.M. MASHAYEKH (2004): Evaluation of the chloroplast DNA among *Vicia faba* L. germplasm using restriction-site analysis. *Iran J Sci Technol* 28(A1): 51-55.
- TABERLET, P., L. GIELLY, G. PATOU, J. BOUVET (1991): Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Mol Biol* 17: 1105-1109.
- TANNO, K., G. WILLCOX (2006): The origins of cultivation of *Cicer arietinum* L. and *Vicia faba* L.: early finds from Tell el-Kerkh, north-west Syria, late 10th millennium B.P. *Veg Hist Archaeobot* 15: 197-204.
- TAMURA, K., D. PETERSON, N. PETERSON, G. STECHER, M. NEI, S. KUMAR (2011): MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28(10): 2731-2739.
- TOLLEFSRUD, M.M., J.H. SØNSTEBØ, C. BROCHMANN, Ø. JOHNSEN, T. SKRØPPA, G.G. VENDRAMIN (2009): Combined analysis of nuclear and mitochondrial markers provide new insight into the genetic structure of North European *Picea abies*. *Heredity* 102(6): 549-562.
- WOLFE, K.H., W.H. LI, P.M. SHARP (1987): Rates of nucleotide substitution vary greatly among plant mitochondrial, chloroplast, and nuclear DNAs. *P Natl Acad Sci USA* 84(24): 9054-9058.

- 
- WOLOSZYNSKA, M. (2010): Heteroplasmy and stoichiometric complexity of plant mitochondrial genomes—though this be madness, yet there's method in't. *J Exp Bot* 61(3): 657-671.
- WON, H., S.S. RENNER (2003): Horizontal gene transfer from flowering plants to *Gnetum*. *P Natl Acad Sci USA* 100(19): 10824-10829.
- VAN DE VEN, W.T.G., N. DUNCAN, G. RAMSAY, M. PHILLIPS, W. POWELL, R. WAUGH (1993): Taxonomic relationships between *V. faba* and its relatives based on nuclear and mitochondrial RFLPs and PCR analysis. *Theor Appl Genet* 86(1): 71-80.
- ZHANG, L.B., M.P. SIMMONS, A. KOCYAN, S.S. RENNER (2006): Phylogeny of the Cucurbitales based on DNA sequences of nine loci from three genomes: implications for morphological and sexual system evolution. *Mol Phylogenet Evol* 39(2): 305-322.

**MOLEKULARNI MARKERI ZA ISTRAŽIVANJA DIVERZITETA NA NIVOU MITOHONDRIJALNOG GENOMA KOD BOBA (*Vicia faba*)**

Jelena M. ALEKSIĆ<sup>1</sup>, Jovanka MILJUŠ-ĐUKIĆ<sup>1</sup>, Živko JOVANOVIĆ<sup>1</sup>, Branko TOMIĆ<sup>2</sup>,  
Bojana BANOVIĆ<sup>1</sup>

<sup>1</sup> Laboratorija za molekularnu biologiju biljaka, Institut za molekularnu genetiku i genetičko inženjerstvo (IMGGE), Univerzitet u Beogradu, Beograd, Srbija

<sup>2</sup> Laboratorija za molekularnu biologiju, Institut za molekularnu genetiku i genetičko inženjerstvo (IMGGE), Univerzitet u Beogradu, Beograd, Srbija

Izvod

Kompletni mitohondrijalni genomi (mtDNK) boba (*Vicia faba*) i dve srodne vrste, *Vigna angularis* i *Vigna radiata*, trenutno dostupni u javnoj banci gena GenBank, su korišćeni za *in silico* PCR analizu kojom je utvrđena primenljivost 15 publikovanih univerzalnih pari prajmera za amplifikuju svih 14 *cis*-isecajućih introna u genima NADH subjedinica (*nad* gena) mitohondrijalnog genoma kod *V. faba* i srodnih vrsta. Zatim je putem PCR amplifikacije analizirano da li su PCR produkti generisani primenom sedam od 15 testiranih pari prajmera kod 16 genotipova *V. faba*, koji su predstavljali sve botaničke varijetete ove vrste (*major*, *minor*, *equina* i subsp. *paucijuga*) različitog stepena oplemenjivanja (tradicionalni i poboljšani kultivari) poreklom iz Evrope, Afrike, Azije i južne Amerike, pogodni za dalju manipulaciju. Dizajnirani su i novi prajmeri za PCR amplifikaciju *nad1* introna 2/3 kod *V. faba* za koji je takođe utvrđena intraspecijska varijabilnost na nivou primarne sekvence nukleotida. Na osnovu rezultata *in silico* predikcija i PCR amplifikacija, definisan je set prajmera za PCR amplifikaciju pet introna u *nad* genima kod *V. faba* koji predstavljaju molekularne markere pogodne za buduće filogeografske i druge studije kod ove vrste za koju ne postoje pouzdani podaci o divljim srođnicima, centru poreklu i domestikaciji.

Primljeno 20. IV. 2014.

Odobreno 05. IX. 2014.